



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>C12N 15/46, 15/62, C07K 14/14, 19/00, A61K 39/15</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/26380</b> <b>(43) International Publication Date:</b> 11 May 2000 (11.05.00)
<b>(21) International Application Number:</b> PCT/US99/25799 <b>(22) International Filing Date:</b> 1 November 1999 (01.11.99) <b>(30) Priority Data:</b> 60/106,347 30 October 1998 (30.10.98) US <b>(71) Applicant:</b> CHILDREN'S HOSPITAL MEDICAL CENTER [US/US]; 3333 Burnet Avenue, Cincinnati, OH 45229-3039 (US). <b>(72) Inventors:</b> CHOI, Anthony; 1055 Lawton Road, Park Hills, KY 41011 (US). WARD, Richard, L.; 2333 Shimmering Bay, Cincinnati, OH 45244 (US). <b>(74) Agent:</b> HUNT, Dale, C.; 16th Floor, 620 Newport Center Drive, Newport Beach, CA 92660 (US).		<b>(81) Designated States:</b> AE, AL, AM, AT, AT (Utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), DM, EE, EE (Utility model), ES, FI, FI (Utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KR (Utility model), KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
<b>(54) Title:</b> ROTAVIRUS SUBUNIT VACCINE  <b>(57) Abstract</b>  <p>The present invention is directed to the generation and use of recombinant rotavirus fusion proteins as immunogens to produce a protective immune response from immunized individuals. In one embodiment, the present invention contemplates a recombinant rotavirus fusion protein vaccine composition comprising a rotavirus subunit protein or immunogenic fragment thereof, and an adjuvant in combination with the recombinant rotavirus subunit fusion protein. In one aspect of this embodiment, the recombinant rotavirus fusion protein comprises a rotavirus subunit protein and a fusion partner protein in genetic association with the rotavirus subunit protein, wherein the fusion partner protein does not interfere with expression and immunogenicity of the rotavirus subunit protein, the fusion partner protein prevents complex formation by the rotavirus subunit protein, and the fusion partner protein facilitates purification of the recombinant rotavirus fusion protein. In another aspect of this embodiment, the rotavirus subunit protein is selected from the group consisting of VP1, VP2, VP3, VP4, VP6, VP7, NSP1, NSP2, NSP3, NSP4 or NSP5. In yet another aspect of this embodiment, the rotavirus subunit protein is VP6.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## ROTAVIRUS SUBUNIT VACCINE

### Background of the Invention

Rotaviruses comprise a genus within the family Reoviridae, and are ubiquitous throughout the animal kingdom. Rotavirus infection is known to cause gastrointestinal disease and is considered the most common cause of gastroenteritis in infants. In fact, essentially every species of domestic animal tested has its own endogenous rotaviruses that cause diarrhea in newborns. Rotaviruses are also thought to be the most significant cause of gastroenteritis in young children and animals. Kapikian & Chanock, *Rotaviruses, Virology*, 2nd edition, Fields *et al.*, eds., New York: Ravenpress, 1353-1404 (1990). Rotaviruses cause diarrheal disease primarily in the young, but infection and disease in older children and adults are also common.

Rotavirus infection is the leading cause of human diarrhea both in developed and developing countries, especially in infants less than one year of age. Worldwide approximately 870,000 children die of rotavirus associated disease every year in developing countries. Even in the United States there are approximately 70 deaths and 200,000 hospitalizations per year due to rotavirus diarrhea. While mortality has been controlled in developed countries by universal access to emergency medical care, morbidity from rotavirus disease remains high.

Around 90% of infants and children develop a rotavirus infection by the third year of life whether they are living in a developed or developing country. The impact of the disease is substantial. An estimated 3,000,000 cases of rotavirus diarrhea occur in the United States annually, leading to some 500,000 physician visits and 55,000-100,000 hospitalizations, which costs the health care system an estimated \$500,000,000 to \$600,000,000 a year. When indirect costs are included, the figure rises to \$1.4 billion. *Pediatrics*, 609-615 (1995).

Rotaviruses are double-stranded RNA viruses and contain a multi-segmented genome. Because the viral genome is arranged in segments, these viruses are capable of genetic reassortment. This reassortment occurs when two or more rotaviruses infect a single cell and the various viral segments reassort during the packaging of new virus particles assembled in the cytoplasm of infected cells. The ability of the virus to reassort its genome components results in a great diversity of immune responses generated against the virus. This ability has also made generation of a rotavirus vaccine extremely challenging.

A variety of different approaches have been taken to generate a rotavirus vaccine suitable to protect human populations from the various serotypes of rotavirus. These approaches include various Jennerian approaches, use of live attenuated viruses, use of virus-like particles, nucleic acid vaccines and viral sub-units as immunogens.

An example of a Jennerian vaccine can be found in Zygraich, U.S. Patent No. 4,341,763, issued July 27, 1982, entitled, "Methods of Vaccinating Humans Against Rotavirus Infection." In this reference, bovine rotaviruses, either attenuated or inactivated, were used to immunize human beings against rotavirus infection. A modified Jennerian approach has also been adopted with the goal of attaining broader antigenic coverage. This approach entailed engineering new reassortant rotavirus types so that a group of viruses would simultaneously display epitopes of various serotypes. However, as with all attenuated viruses, there is always a high risk of reversion.

The Food and Drug Administration has recently approved a multivalent vaccine against rotavirus making it available to pediatricians in the United States. The vaccine, which was developed by Albert Z. Kapikian, MD, and colleagues at the Laboratory of Infectious Diseases (NIAID) is an excellent example of a multivalent vaccine. A recent reported study conducted in Caracas testing this vaccine was reported in the New England Journal of Medicine 1997; 337, 1181-1187. Among some 2,200 infants enrolled in the double-blind, placebo controlled study, the vaccine was successful in reducing severe diarrheal illness in about 70% of subjects but was less effective at reducing the incidence of less serious diarrheal illness. Recently, it was reported that there have been serious side effects such as bowel blockage associated with this vaccine.

Another example of the modified Jennerian approach is found in Clark *et al.*, U.S. Patent No. 5,626,851, issued May 6, 1997, entitled, "Rotavirus Reassortant Vaccine." In this example a rotavirus reassortant suitable for use as a vaccine was produced using genetic reassortment between an attenuated bovine rotavirus and at least one rotavirus representing an epidemiologically important serotype. This reference looked to create a bovine rotavirus that carried either VP4 or VP7 genes which, when presented to a immunologically naive subject, would induce a protective immune response therein. Thus, this method relies upon the use of whole viruses to immunize a subject.

U.S. Patent Nos. 4,624,850, 4,636,385, 4,704,275, 4,751,080, 4,927,628, 5,474,773, and 5,695,767, each describe a variety of rotavirus vaccines and/or methods of preparing the same. A commonality shared by the members of this group is that each of these vaccines relies on the use of whole viral particles to create the ultimate rotavirus vaccines. Given the long standing need for an effective, multivalent vaccine, it is clear that this body of work has been only partially successful in addressing the need for such a vaccine.

Departing from traditional methods of vaccine generation, advances in the field of molecular biology have permitted the expression of individual rotavirus proteins. Using these techniques, vaccine candidates generated from virus-like particles of different protein compositions have shown potential as subunit vaccines. In one reference, VLPs containing VPs 2 and 6 or VPs 2, 6, and 7 were administered to mice with and without the addition of cholera toxin. O'Neal *et al.*, "Rotavirus Virus-like Particles Administered Mucosally Induce Protective Immunity," *J. Virology*, 71(11):8707-8717 (1997). Both types of VLPs induced protective immunity in immunized mice, although protection was more effective when the VLPs were administered with cholera toxin (CT). In a subsequent study by the same group, the *Escherichia coli* heat-labile toxin (LT) was compared to CT for effectiveness in producing rotavirus protection. O'Neal *et al.*, "Rotavirus 2/6 Virus-like Particles Administered Intranasally with Cholera Toxin, *Escherichia coli* Heat-Labile Toxin (LT), and LT-R192G Induce Protection from Rotavirus Challenge," *J. Virology*, 72(4):3390-3393 (1998). This group concluded that both the wild-type LT and a recombinant form of the molecule were effective adjuvants when immunizing with rotavirus VLPs.

Core-like particles and VLPs have also been used to immunize cows. Fernandez, *et al.*, "Passive Immunity to Bovine Rotavirus in Newborn Calves Fed Colostrum Supplements From Cows Immunized with Recombinant SA11 rotavirus core-like particle (CLP) or virus-like particle (VLP) vaccines," *Vaccine*, 16(5):507-516 (1998). In this study the

ability of CLPs and VLPs to create passive immunity was studied. This group concluded that VLPs were more effective than CLPs in inducing passive immunity.

In several studies the individual viral proteins have also been used to immunize subjects. For example, one group used baculovirus-expressed VP4 protein from the simian rhesus rotavirus (RRV) to parenterally immunize murine dams. Newborn mice suckling from immunized dams were found to be protected against RRV challenge and against challenge by a murine rotavirus through a passive immunity scheme. Mackow *et al.*, "Immunization with baculovirus-Expressed VP4 Protein Passively Protects Against Simian and Murine Rotavirus Challenge," *J. Virol.* 64(4): 1698-1703 (1990).

Rotavirus proteins have even been used as immunological carrier complexes to facilitate the presentation of other epitopes to a subject. In one reference, VP6 was chosen as a carrier molecule for a particular antigen, based on the viral protein's ability to bind peptides. Sabara, *et al.*, U.S. Patent No. 5,374,426, issued December 20, 1994, entitled, "Rotavirus Nucleocapsid Protein VP6 in Vaccine Compositions."

Exploiting another avenue, research has also been performed where a protective immune response was elicited using a DNA vaccine. Herrmann, *et al.*, U.S. Patent No. 5,620, 896, issued April 15, 1997, entitled, "DNA Vaccines Against Rotavirus Infections." While the results from this method are interesting, the degree of protection found in mice immunized with plasmids containing either the VP4, VP7 or VP6 genes in a murine retrovirus were very limited. Moreover, there are at least two reports of rotavirus DNA vaccines that failed to provide any protection to the immunized animal from rotaviral infection. Choi, *et al.*, "Particle Bombardment-Mediated DNA Vaccination with Rotavirus VP6 Induces High Levels of Serum Rotavirus IgG but Fails to Protect Mice Against Challenge," *Virology* 232: 129-138 (1997). Choi *et al.* "Particle Bombardment-Mediated DNA Vaccination With Rotavirus VP4 or VP7 Induces High Levels of Serum IgG but Fails to Protect Mice Against Challenge," *Virology* 250:230-240 (1998).

This review of the state of the art attempts to provide some measure of the extent of time and energy that has been expended to date to develop an effective rotavirus vaccine. Yet, even with the most successful efforts to date, even the severe forms of rotaviral disease can be prevented only about 75% of the time. Given this limitation, there remains a need for a safe and effective rotavirus vaccine, which overcomes the above deficiencies. The vaccine of present invention is designed to remedy and eliminate these problems.

#### Summary of the Invention

The invention disclosed herein relates to compositions comprising various rotaviral proteins and methods of using these compositions to provide protection against rotaviral disease. One embodiment of the invention is a composition comprising a rotavirus VP6 protein or a fragment thereof, and an adjuvant in a pharmaceutical carrier, wherein said adjuvant is effective in generating a disease-reducing response to said VP6 protein.

Another embodiment of the invention encompasses a recombinant rotavirus fusion protein composition, comprising: a rotavirus subunit fusion protein or fragment thereof, a fusion protein partner in genetic association with said recombinant rotavirus subunit protein or fragment thereof, and an adjuvant in a pharmaceutical carrier, wherein said adjuvant is effective in stimulating a disease-reducing immunogenic response to said rotavirus fusion protein.

A full-length DNA copy of a gene encoding a recombinant rotavirus fusion protein, wherein said gene encoding said recombinant rotavirus protein comprising a rotavirus subunit protein or an immunogenic fragment thereof, and a fusion partner protein is contemplated in another embodiment of the invention disclosed herein.

5 A host cell comprising a DNA clone encoding recombinant rotavirus proteins is contemplated in another embodiment of the disclosed invention.

The invention disclosed also contemplates a computer readable medium having recorded thereon peptide sequences of rotavirus proteins selected from the group consisting of SEQ ID NOs:15-25.

10 A number of methods are disclosed as part of the invention. For example, one embodiment discloses a method of generating an immune response in a subject in need of rotavirus immunity comprising the steps of administering an immunogenic composition of comprising a rotavirus protein or fragment thereof, and an adjuvant in a pharmaceutically acceptable form to said subject and generating a disease-reducing immunogenic response in said subject.

15 Another embodiment of the invention is a method of immunizing a subject in need of rotavirus immunity comprising the steps of: immunizing with a rotavirus vaccine; and subsequently immunizing said subject with a recombinant rotavirus vaccine composition, whereby immunological protection of said individual against rotavirus infection is increased.

#### Brief Description of the Drawings

20 FIGURE 1 shows some of the important features of pMAL-c2. Using this plasmid, recombinant plasmids were constructed which express chimeric proteins containing, the entire VP6, portions of VP6, the entire VP4 or a truncated form of VP7. pMAL-c2 contains a promoter sequence called *P<sub>tac</sub>* which controls transcription of the fusion gene *malE-lac Z $\alpha$* . The gene *malE-lac Z $\alpha$*  encodes a chimeric protein containing MBP and the  $\alpha$  fragment of the enzyme  $\beta$ -galactosidase. Rotavirus gene sequences were cloned into *Xmn* I, which is one of the multiple restriction sites present in the plasmid used for gene cloning. The created recombinant plasmids express chimeric proteins containing rotavirus proteins that are genetically fused with MBP. The precise site for *Xmn* I insertion is indicated by the arrow in FIGURE

25 1. The same arrow also marks the exact position in the protein which can be enzymatically digested by Factor Xa in some, but not all, chimeric MBP proteins. pMAL-c2 also contains the *lac I'* sequence which encodes a repressor protein that suppresses *P<sub>tac</sub>* controlled transcription until IPTG is added. The plasmid also contains the *colE1* origin of replication in *E. coli* and the ampicillin resistance gene that are typical features of many bacterial plasmids.

30 FIGURE 2 shows the results of Western blot analysis of chimeric MBP proteins, which contain MBP genetically fused with various rotavirus proteins. Aliquots of the purified proteins were subjected to separation by SDS-PAGE. Western blot analyses were then carried out by blotting separated proteins onto nitrocellulose sheets. Antibodies generated against MBP were used and a chromogenic reaction was performed to visualize chimeric proteins derived from unfractionated cells (whole cell lysate) before the first step and purified proteins after the last step of the purification scheme. The arrows indicate the chimeric proteins containing MBP genetically fused with VP4, VP6 or truncated VP7.

35

FIGURE 3 shows evidence to establish that chimeric MBP::VP6 protein does not form structures that resemble rotavirus-like particles. Purified MBP::VP6 was put on the top of a sucrose gradient which was layered on top of a cesium chloride cushion. Rotavirus particles that were devoid of VP4 and VP7 were also layered on an identical sucrose gradient/cesium chloride cushion. By subjecting the protein and virus particles to a centrifugal force, the majority of the rotavirus particles traversed into fraction number 11 and 12 of the sucrose gradient and into fraction number 16 containing cesium chloride. In contrast, chimeric MBP::VP6 did not traverse into the gradient but remained in the top fractions (number 1 and 2). These results clearly demonstrated that the presence of MBP in the fusion protein does not allow MBP::VP6 to form any structures that resemble rotavirus particles.

FIGURE 4. Rotavirus shedding in BALB/c immunized with MBP::VP6 and LT (attenuated *E. coli* heat labile toxin) or LT only following challenge with EDIM. Groups of 8 BALB/c mice were intranasally immunized with two doses, separated by 14 days, of either 8.8ug MBP::VP6 and 10 µg of LT or 10 µg of LT alone. Four weeks after the last immunization, the mice were challenged with  $4 \times 10^4$  focus forming units (ffu) of EDIM (P9 12/15/97). Stools were collected for seven days following challenge and were tested for rotavirus antigen by enzyme-linked immunosorbent assay (EIA). The  $A_{490}$  values are the averages obtained each day.

FIGURE 5 illustrates the region of VP6, which may assist in creating minimal subunit rotavirus vaccines. VP6, which consist of 397 amino acid residues, was delineated into four regions: regions A, B, C and D. The exact amino residues delineated by these regions were indicated. The gene sequences that encode regions A and B, B and C, or C and D were cloned into pMAL-c2 at the *Xmn* I site. The plasmids were given the names pMAL-c2/EDIM<sub>AB</sub>, pMAL-c2/EDIM<sub>BC</sub> and pMAL-c2/EDIM<sub>CD</sub> respectively.

FIGURE 6 provides evidence that mice inoculated with chimeric MBP proteins containing regions A and B or containing regions C and D generated IgG which specifically recognized VP6. In these experiments, rotavirus particles were subjected to SDS-PAGE. The separated proteins were then transferred to nitrocellulose sheets. The sheets were cut into strips. Individual strips were incubated with a specific immune serum sample collected from mice inoculated with pMAL-c2/EDIM<sub>AB</sub>, pMAL-c2/EDIM<sub>BC</sub> or pMAL-c2/EDIM<sub>CD</sub>. While pMAL-c2/EDIM<sub>AB</sub> and pMAL-c2/EDIM<sub>CD</sub>-inoculated mice generated anti-VP6 IgG, no specific IgG was detected by pMAL-c2/EDIM<sub>BC</sub>-immunized mice.

FIGURE 7 further defines the CD region of VP6 in order to determine whether an even smaller minimal subunit rotavirus vaccine can be attained. The CD region was delineated into four regions: regions 1, 2, 3 and 4. The exact amino residues delineated by these regions were indicated. Recombinant plasmids were constructed using pMAL-c2 to harbor regions 1, 2, 3, and 4. The plasmids were given the names pMAL-c2/EDIM<sub>CD1</sub>, pMAL-c2/EDIM<sub>CD2</sub>, pMAL-c2/EDIM<sub>CD3</sub> and pMAL-c2/EDIM<sub>CD4</sub> respectively.

#### Detailed Description of the Preferred Embodiment

This invention relates to the generation of rotavirus subunit proteins for use in vaccines and methods of providing protective immunity to vertebrates, including humans, against rotavirus infection or disease. As one embodiment, the protective immunity generated by vaccines containing the recombinant rotavirus proteins of the present invention is a dominantly cell-mediated immune response. This immune response may interfere with the

infectivity or activity of the rotavirus, or it may limit the spread or reproduction of the virus. The immune response resulting from vaccination with a vaccine containing the proteins of the present invention provides protection against subsequent challenge by a homologous or heterologous rotavirus.

5 The vaccines of the present invention are composed of a native recombinant rotavirus protein or immunogenic fragment(s) thereof, a rotavirus fusion protein, or immunogenic fragment(s) thereof, an adjuvant, and a pharmaceutically acceptable carrier. According to one embodiment of the present invention, a composition comprising a rotavirus protein or an immunogenic portion thereof is genetically associated with a fusion protein partner, and an adjuvant such as the A1 subunit, the B subunit of cholera toxin or *E. coli* heat-labile toxin present in a pharmaceutically acceptable carrier. This composition is administered to an individual in whom an immune response directed against the  
10 rotavirus subunit protein is sought and protection against rotavirus infection and disease is desired.

The rotavirus native recombinant, or fusion proteins of the present invention may be composed of any rotavirus protein product or immunogenic fragment thereof. The rotavirus protein may be chosen from any of the structural or non-structural viral proteins encoded by the rotavirus genome. For example, in one embodiment, the rotavirus protein or immunogenic fragment thereof may be chosen by selecting from the group of rotavirus genome  
15 segments consisting of segment 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or 11, that encode rotavirus protein sequences. In another embodiment, the rotavirus protein or immunogenic fragment thereof may be selected from the group of rotavirus structural proteins consisting of VP1, VP2, VP3, VP4, VP6 and VP7. In another embodiment, the rotavirus protein may be selected from the group of rotavirus non-structural proteins consisting of NSP1, NSP2, NSP3, NSP4 and NSP5. In yet another embodiment, the rotavirus protein used in the fusion protein construct is VP6. In still  
20 another embodiment, the use of an immunogenic fragment of VP6 may be used in the present invention.

The rotavirus recombinant native or fusion proteins of the present invention may be used in a vaccine composition at a concentration effective to elicit an immune response from an immunized subject. The concentration of rotavirus proteins of the present invention may range from about 0.01 µg/ml to 1 mg/ml. In another embodiment, the concentration of rotavirus proteins used in a vaccine composition may range from about 0.1 µg/ml to 100 µg/ml.  
25 In yet another embodiment, the concentration of rotavirus proteins used in a vaccine composition may range from about 1.0 µg/ml to 10 µg/ml. In still another embodiment, the concentration of rotavirus proteins used in a vaccine composition may be about 8.8 µg/ml. These ranges are provided for the sake of guidance in practicing the present invention. It should be noted that other effective concentrations of recombinant rotavirus proteins may be determined by one of ordinary skill in the art using experimental techniques well known in that art.

30 The rotavirus fusion proteins contemplated by the present invention are composed of a suitable fusion protein partner in genetic association with a rotavirus protein or immunogenic fragment thereof. The term in genetic association refers to a contiguous sequence of amino acids produced from a mRNA produced from a gene containing codons for the amino acids of the rotavirus protein and the fusion protein partner. A suitable fusion protein partner consists of a protein that will either enhance or at least not diminish the recombinant expression of the rotavirus fusion  
35 protein product when the two are in genetic association. Further, a suitable fusion protein partner may actively



prevent the assembly of the rotavirus fusion proteins into multimeric forms after the rotavirus fusion protein has been expressed. For example, the fusion protein partner should prevent the formation of dimers, trimers or virus-like structures that might spontaneously form if the rotavirus protein were recombinantly expressed in the absence of the fusion protein partner. Still further, a suitable fusion partner will facilitate the purification of the chimeric rotavirus fusion protein. A representative list of suitable fusion protein partners includes maltose binding protein, poly-histidine segments capable of binding metal ions, inteine, antigens to which antibodies bind, S-Tag, glutathione-S-transferase, thioredoxin,  $\beta$ -galactosidase, nonapeptide epitope tag from influenza hemagglutinin, a 11-amino acid epitope tag from vesicular stomatitis virus, a 12-amino acid epitope from the heavy chain of human Protein C, green fluorescent protein, cholera holo toxin or its B subunit, *E. coli* heat-labile holotoxin or its B subunit, CTA1-DD, streptavidin and dihydrofolate reductase.

The invention is also directed toward producing rotavirus proteins for use in vaccines directed to protect immunized individuals from rotavirus infection and/or disease. Accordingly, the invention contemplates the use of an adjuvant, such as an immunogenic protein, effective to induce desirable immune responses from an immunized animal. Such a protein must possess those biochemical characteristics required to facilitate the induction of a protective immune response from immunized vertebrates while simultaneously avoiding toxic effects to the immunized animal.

In one embodiment of the present invention, rotavirus recombinant native or fusion proteins are mixed with an adjuvant such as a bacterial toxin. The bacterial toxin may be a cholera toxin. Alternatively, the rotavirus fusion protein may be mixed with the B subunit of cholera toxin (CTB). In another embodiment, an *E. coli* toxin may be mixed with the rotavirus fusion protein. For example, the rotavirus fusion protein may be mixed with *E. coli* heat-labile toxin (LT). The rotavirus fusion proteins of the present invention may be mixed with the B subunit of *E. coli* heat-labile toxin (LTB) to form a vaccine composition. Other adjuvants such as cholera toxin, labile toxin, tetanus toxin or toxoid, poly[di(carboxylatophenoxy)phosphazene] (PCPP), saponins Quil A, QS-7, and QS-21, RIBI (HAMILTON, MT), monophosphoryl lipid A, immunostimulating complexes (ISCOM), Syntax, Titer Max, M59, CpG, dsRNA, and CTA1-DD (the cholera toxin A1 subunit (CTA1) fused to a dimer of the Ig-binding D-region of *Staphylococcus aureus* protein A (DD)), are also contemplated.

The adjuvants discussed above may be used in a vaccine composition at a concentration effective to assist in the eliciting of an immune response against the recombinant rotavirus fusion proteins of the present invention from an immunized subject. The concentration of adjuvant included in the vaccine compositions of the present invention may range from about 0.01  $\mu\text{g/ml}$  to 1 mg/ml. In another embodiment, the concentration of adjuvant used in a vaccine composition may range from about 0.1  $\mu\text{g/ml}$  to 100  $\mu\text{g/ml}$ . In yet another embodiment, the concentration of adjuvant used in a vaccine composition may range from about 1.0  $\mu\text{g/ml}$  to 100  $\mu\text{g/ml}$ . In still another embodiment, the concentration of adjuvant used in a vaccine composition may be about 10.0  $\mu\text{g/ml}$ . These ranges are provided for the sake of guidance in practicing the present invention. It should be noted that other effective concentrations of adjuvants may be determined by one of ordinary skill in the art using experimental techniques well known in that art.

The invention also contemplates immunization with a rotavirus fusion protein, a recombinant native protein, or a fragment or fusion fragment, and a suitable adjuvant contained in a pharmaceutically acceptable composition. Such a composition should be sterile, isotonic, and provide a non-destabilizing environment for the rotavirus fusion protein and the adjuvant. Examples of this are buffers, tissue culture media, various transport media and solutions containing proteins (such as BSA), sugars (sucrose) or polysaccharides.

The vaccine compositions of the invention contain conventional pharmaceutical carriers. Suitable carriers are well known to those of skill in the art. These vaccine compositions may be prepared in liquid unit dose forms. Other optional components, *e.g.*, stabilizers, buffers, preservatives, excipients and the like may be readily selected by one of skill in the art. However, the compositions may be lyophilized and reconstituted by the individual administering the vaccine prior to administration of the dose. Alternatively, the vaccine compositions may be prepared in any manner appropriate for the chosen mode of administration, *e.g.*, intranasal administration, oral administration, etc. The preparation of a pharmaceutically acceptable vaccine, having due regard to pH, isotonicity, stability and the like, is within the skill of the art.

The dosage regimen involved in a method for vaccination, including the timing, number and amounts of booster vaccines, will be determined considering various hosts and environmental factors, *e.g.*, the age of the patient, time of administration and the geographical location and environment.

Also included in the present invention are methods of vaccinating humans against rotavirus infection and disease with the novel rotaviral proteins and vaccine compositions described above. The vaccine compositions, comprising a full-length rotavirus protein, a rotavirus fusion protein, a recombinant native protein or fragments and fusion fragments, mixtures of the above, and an adjuvant described herein may be administered by a variety of routes contemplated by the present invention. Such routes include intranasal, oral, rectal, vaginal, intramuscular, intradermal and subcutaneous administration.

Vaccine compositions for parenteral administration include sterile aqueous or non-aqueous solutions, suspensions or emulsions, the protein vaccine, and an adjuvant as described herein. The composition may be in the form of a liquid, a slurry, or a sterile solid which can be dissolved in a sterile injectable medium before use. The parenteral administration is preferably intramuscular. Intramuscular inoculation involves injection via a syringe into the muscle. This injection can be via a syringe or comparable means. The vaccine composition may contain a pharmaceutically acceptable carrier. Alternatively, the present vaccine compositions may be administered via a mucosal route, in a suitable dose, and in a liquid form. For oral administration, the vaccine composition can be administered in liquid, or solid form with a suitable carrier.

Doses of the vaccine compositions may be administered based on the relationship between the concentration of the rotavirus fusion protein contained in the vaccine composition and that concentration of fusion protein required to elicit an immune response from an immunized host. The calculation of appropriate doses to elicit a protective immune response using the rotavirus fusion protein vaccine compositions of the present invention are well known to those of skill in the art.

A variety of immunization methods are contemplated by the invention to maximize the efficacy of the rotavirus protein vaccine compositions described herein. In one embodiment, females of offspring-bearing age are immunized with the vaccines of the invention. In this embodiment, immunized females develop a protective immune response directed against rotavirus infection or disease and then passively communicate this protection to an offspring by nursing. In another embodiment, newborns are immunized with the vaccine compositions of the invention and shortly thereafter the nursing mother is immunized with the same vaccine. This two tiered approach to vaccination provides the newborn with immediate exposure to viral epitopes that may themselves be protecting. Nevertheless, the passive immunity supplied by the mother would augment the protection enjoyed by the offspring. This method would therefore provide the offspring with both active and passive protection against rotavirus infection or disease.

In still another embodiment, an individual is immunized with the vaccine composition of the invention subsequent to immunization with a multivalent vaccine. The immunization of a subject with two different vaccines may synergistically act to increase the protection an immunized individual would enjoy over that obtained with only one vaccine formulation. One draw back of immunization with a multivalent live virus vaccine formulation is that booster immunizations with the same vaccine are usually not effective. In this embodiment of the invention, the vaccine compositions serve as such a booster to increase the protection of the immunized individual against rotaviral infection or disease.

The following examples teach the generation of all types of rotavirus protein vaccine compositions. These examples are illustrative and are not intended to limit the scope of the present invention. One of skill in the relevant art would be able to use the teachings described in the following examples to practice to full scope of the present invention.

## EXAMPLES

### Example 1

#### Construction of recombinant pMAL-c2 plasmids

Recombinant plasmids pMAL-c2/EDIM4, pMAL-c2/EDIM6 and pMAL-c2/LDIM7 were constructed using pMAL-c2 (New England Biolabs, Beverly MA) by insertion of cDNAs encoding full length VP4 or VP6, or a truncated form of VP7 (TrVP7) of rotavirus strain EDIM (FIGURE 1). cDNAs were synthesized by polymerase chain reaction (PCR) using the plasmids pGEM-3Z/EDIM4, pGEM-3Z/EDIM6 and pGEM-3Z/EDIM7 as templates and gene specific primers determined by nucleotide sequencing of the gene inserts. The nucleotide sequences have been deposited into GenBank nucleotide sequence database and assigned with the Accession Numbers AF039219, U65988 and AF039220 for VP4, VP6 and VP7 gene respectively.

The murine EDIM strain of rotavirus used for the construction of the pGEM recombinant plasmids was originally isolated from the stool of an infected mouse and adapted to grow in cell culture by passage in MA-104 cells in the laboratory. A triply plaque-purified isolate of the ninth passage was used to infect MA-104 cells to yield stock virus for RNA purification. To generate cDNAs of rotavirus genes encoding strain VP4, VP6 and VP7, reverse transcription/polymerase chain reaction was carried out using purified genomic rotavirus RNA, a forward and a reverse

primer obtained from the untranslatable regions of the gene. The cDNAs generated by RT/PCR were cloned into the *Sma* I site of the multiple cloning site of pGEM-3Z (Promega, Madison, WI). Ligation products were then transformed into *E. coli*. White transformants carrying recombinant plasmids were selected by growing cells on LB agar plates containing IPTG (0.5 mM) and X-gal. Plasmids from individual colonies were purified and were analyzed by nucleotide sequencing.

The cDNAs generated by PCR were inserted into the restriction site *Xmn* I of pMAL-c2, placing the inserted sequences downstream from and in genetic association with the *E. coli* malE gene, which encodes maltose binding protein (MBP), resulting in the expression of MBP fusion protein. The plasmid utilized the strong "tac" promoter and the malE translation initiation signals to give high-level expression of the fusion protein. pMAL-c2 contains the factor Xa cleavage site that is located downstream from the malE sequence to enable cleavage of the heterologous protein from MBP. The plasmid conveyed ampicillin resistance to recombinant bacteria and a lacZ-alpha gene sequence for blue-to-white selection of recombinants with inserts.

Following ligation of cDNA and *Xmn*I-digested pMAL-c2, recombinant pMAL-c2 plasmids were transformed into *E. coli*. White colonies of bacteria containing recombinant plasmids on an agar plate were then identified in the presence of IPTG and X-gal, and selected for further screening by PCR for gene identity and orientation. Nucleotide sequencing was used to ultimately confirm the authenticity of the rotavirus gene sequence.

#### Example 2

#### Expression and Purification of Fusion Proteins

Recombinant bacteria were grown as an overnight culture (37°C, shaken at 215 rpm) in rich broth (tryptone, 10 gm; yeast extract, 5 NaCl, 5 gm; glucose, 2 gm and 100 mg of ampicillin per liter). On the following day, 10 ml of overnight cell culture were inoculated into 1 liter of rich broth containing glucose and ampicillin. The culture was grown until the optical density  $A_{600}$  reached 0.6. IPTG was then added to 0.3 mM to induce expression of fusion protein. Growth was continued for 3 hours.

Cells were harvested by centrifugation (4,000 g; 20 min at 4°C), resuspended in PBS, and subjected to centrifugation. The pellet was frozen at -20°C, thawed slowly in cold water, and resuspended in a total of 50 ml of buffer L (5 mM  $\text{NaH}_2\text{PO}_4$ , 10 mM  $\text{Na}_2\text{HPO}_4$ , 30 mM NaCl, 10 mM 2-beta mercaptoethanol and 0.2% Tween 20, 1 mM PMSF, 25 mM benzamidine, and 200 mg/L of lysozyme). After digestion for 15 min at room temperature (rt), the suspension was sonicated by three 30 second bursts (BioSonic IV, 50% power setting) while placed in an ice/water bath. NaCl (26.5 mg/ml) and RNase A (5  $\mu$ l of 10 mg/ml) were added to each 10 ml of sonicate which was then centrifuged (54,000 g, 30 min) to obtain a supernatant containing a crude preparation of fusion protein.

Fusion proteins in the crude preparation were purified by affinity chromatography. Amylose resin (New England Biolab, Beverly MA) was prepared by placing 25 ml of the packed resin in a 250 ml centrifuge tube and washed twice with eight volumes of buffer C (Buffer L containing 0.5 M NaCl). For each wash, the mixture was rocked for 30 min at 4°C, and the resin was recovered by centrifugation (2,100 g, 5 min). The supernatants, which contained the fusion proteins, were mixed with amylose resin for 2 hours in a 500 ml flask on a magnetic stirrer. After centrifugation (2,100 g, 5 min), the resin was recovered, then resuspended in 50 ml of buffer C, rocked for 30 min and

finally centrifuged to recover the resin. The resin was washed in this manner for a total of 3 times and finally washed overnight with 500 ml of buffer C.

On the following day, the resin was recovered by centrifugation (2,100 *g*, 5 min) and resuspended in 50 ml of buffer D (50 mM Tris-HCl, pH 7.5; 50 mM NaCl; 1 mM EDTA; 10 mM 2-beta mercaptoethanol; 1 mM PMSF), and rocked for 30 min. The resin was spun down and the bound fusion proteins were eluted from the resin with 250 ml of 15 mM maltose in buffer D for 2 hours. The resin was recovered by centrifugation (2,100 *g*, 5 min) and the supernatant containing the fusion proteins was subjected to buffer exchange to PBS and was simultaneously concentrated by ultrafiltration using a stirred-cell concentrator (Amicon, Beverly MA; model 8400). The purified fusion proteins were analyzed by Western blot analyses (FIGURE 2).

### Example 3

#### Biochemical characterization of MBP::VP6 fusion protein

It has been shown that recombinant VP6 expressed by the baculovirus expression system forms structures that resemble double-layered rotavirus particles when examined by electron microscopy. Purified MBP::VP6 fusion protein was analyzed by sucrose gradients to determine if these fusion proteins assembled into organized structures resembling virus particles that could be fractionated in a sucrose gradient. MBP::VP6 was subjected to centrifugation (SW 50, 35,000 *g*, 60 min) through a 4 ml sucrose gradient (20-50%) on a 1 ml cesium chloride cushion (60%). A total of 16, 300- $\mu$ l fractions were collected. Distribution of MBP::VP6 in the sucrose gradient and cesium chloride cushion was analyzed by Western blot analysis and distribution of virus particles was analyzed by silver nitrate staining of the SDS-gel (FIGURE 3). The results showed that MBP::VP6 remained in the top 4 fractions of the gradient, while double-layered virus particles devoid of VP4 and VP7 were recovered from fraction #11 to #12 of the sucrose gradient and in the cesium chloride cushion (fraction #16). The difference in the distribution behavior of MBP::VP6 in the gradient indicated that the fusion protein does not form virus-like structures.

#### Analysis of the Immune Response from Subjects Immunized with

#### VP-6 Fusion Proteins

### Example 4

#### Method of Vaccination and Challenge

Six-week-old virus antibody free female BALB/c mice were purchased from Harlan Sprague-Dawley (Indianapolis, IN). Animals were housed four animals to a cage in sterile micro barrier cages. Between four and ten animals were included in each group. Animals were ear tagged and a blood and stool specimen was collected from each animal prior to vaccination.

Expressed fusion protein of EDIM VP6 or portions of VP6 were used as the immunizing antigens. Protein concentration was calculated to be 176 ng/ $\mu$ l. Animals received 50  $\mu$ l of VP6 (8.8  $\mu$ g) per dose. Animals received either two or three doses separated by two-week intervals. The adjuvant used was *E. coli* LT (R192G) at 1 mg/ml received from Dr. John Clements (Tulane University). The LT was resuspended in deionized H<sub>2</sub>O and 10 mM CaCl<sub>2</sub>. Intranasal inoculations included 10  $\mu$ g LT with antigen. Adjuvant and antigens were mixed prior to immunization.

WO 00/26380

Animals were immunized intranasally (i.n.) by lightly anesthetizing with metofane and instilling approximately 5  $\mu$ l per nostril until the entire dose was delivered.

Four weeks after the last immunization, animals were bled and a stool specimen was collected from each animal to measure antibody responses. Animals were challenged with 100  $\mu$ l of a 1:25 fold dilution of EDIM P9 12/15/97 1 X 10<sup>7</sup> ffu/ml to give a dose of 4 X 10<sup>4</sup> ffu or 100 ID<sub>50</sub>. Stool specimens [two pellets in 0.5 ml of Earl's Balanced Salt Solution (EBSS)] were collected from each mouse for seven days and stored at -20°C. Rotavirus antigen was measured in the stools by EIA to determine shedding.

Twenty-one days after challenge, sera and stool specimens were obtained again to measure antibody responses.

#### Example 5

##### EIA Method to Measure Rotavirus Antigen in Stool to Determine Shedding

Stool specimens for Example 4 were thawed, homogenized and centrifuged (500 g, 10 min). For rotavirus antigen determination, 96-well EIA plates (Corning Costar Co., Corning, NY) were coated overnight at 4°C with 100  $\mu$ l per well of either rabbit antibody to rotavirus (duplicate positive wells) or preimmune rabbit serum (duplicate negative wells). Plates were washed and 50  $\mu$ l of stool supernatant was added to duplicate wells coated with each antibody. After one hour incubation at 37°C on a rotation platform, plates were washed and 50  $\mu$ l normal goat serum (Vector Laboratory, Inc., Burlingame, CA) diluted 100-fold in phosphate-buffered saline containing 5% nonfat dry milk (PBS-M) was added for 15 minutes at room temperature. Fifty microliters of guinea pig antibody to rotavirus diluted 1:500 in PBS-M containing a 1:50 dilution of normal rabbit serum (DAKO, Carpinteria, CA) was added and incubated for 30 minutes. Plates were washed and 50  $\mu$ l of a 1:200 dilution of biotinylated goat anti-guinea pig IgG (Vector) in PBS-M containing a 1:50 dilution of normal rabbit serum was added and incubated 30 minutes. After washing plates, 50  $\mu$ l of a 1:100 dilution of peroxidase-conjugated avidin-biotin (Vector) in wash buffer was added and incubated 30 minutes. The plates were washed and 50  $\mu$ l substrate (*o*-phenylenediamine with H<sub>2</sub>O<sub>2</sub> in citric acid-phosphate buffer) was added and incubated (room temperature) for 15 minutes. The reaction was stopped with 75  $\mu$ l of 1.0 M H<sub>2</sub>SO<sub>4</sub>. The absorbance at 490 nm was measured and the net optical densities were determined by subtracting the average of the negative wells from the average of the positive wells. The specimen was considered positive for rotavirus if the average absorbance of the positive wells was greater than or equal to two times that of the negative wells and greater than or equal to 0.15.

A time course of fecal shedding of rotavirus in mice challenged with EDIM is shown in FIGURE 4. The open circles represent data points from EDIM challenged control mice exposed to a control vaccination containing the LT adjuvant but lacking the rotavirus fusion proteins of the present invention. The filled squares represent data points from EDIM challenged experimental mice that were intranasally vaccinated with the VP6::MBP rotavirus fusion vaccine composition of the present invention. As can be seen from the figure, the incidence of fecal shedding increased from the first day after EDIM challenge in the control mice until reaching a maximum value on the fourth day after challenge. In contrast, mice vaccinated with the VP6::MBP rotavirus fusion protein vaccine composition produce little fecal

shedding over the same period of time. These data clearly show that intranasal vaccination of mice with a VP6::MBP rotavirus fusion vaccine composition greatly reduced the incidence of fecal shedding of virus after rotavirus EDIM challenge. Correlating levels of fecal shedding to levels of infection, vaccination with the VP6::MBP rotavirus fusion vaccine composition of the present invention provides protection against EDIM challenge in mice.

#### Example 6

##### EIA Method to Measure Serum Rotavirus IgG and IgA and Stool Rotavirus IgA

Serum rotavirus IgA and IgG and rotavirus stool IgA were measured as follows. EIA plates (Corning Costar Co., Corning, NY) were coated overnight at 4°C with anti-rotavirus rabbit IgG. After washing with phosphate buffered saline plus 0.05% Tween 20, 50 µl of EDIM viral lysate or mock-infected cell lysate were each added to duplicate positive and duplicate negative wells and plates were incubated for one hour at 37°C on a rotation platform. After washing plates, 50 µl of serial two-fold dilutions of pooled sera from EDIM infected mice assigned concentrations of 160,000 or 10,000 units/ml of rotavirus IgG or IgA, respectively, were added to duplicate wells coated with either EDIM-infected or uninfected MA104 cell lysates to generate a standard curve. Serial 10-fold dilutions of mouse sera to be tested were also added to duplicate wells of each lysate and incubated 1 hour. This was followed by sequential addition of biotin-conjugated goat anti-mouse IgG or IgA (Sigma Chemical Co., St. Louis, MO), peroxidase-conjugated avidin-biotin (Vector Laboratories), and *o*-phenylenediamine substrate (Sigma Chemical Co.). Color development was stopped after fifteen minutes with 1 M H<sub>2</sub>SO<sub>4</sub> and the A<sub>490</sub> was measured. Titers of rotavirus IgG or IgA, expressed as units/ml, were determined from the standard curve generated by subtraction of the average A<sub>490</sub> values of the duplicate cell lysate wells from the average of the wells coated with EDIM lysate.

For determination of stool rotavirus IgA, two stool pellets were collected into 0.5 ml of EBSS, homogenized, and centrifuged (1,500 *g*, 5 min). Stool rotavirus IgA was then measured by the method described above.

The data from these experiments showed that mice immunized with the recombinant MBP::VP6 rotavirus fusion protein vaccines generated an immune response directed against the VP6 fusion protein. Both serum IgG and IgA responses were noted. The serum IgG responses were higher than those of the IgA responses. Concerning the IgG responses, administration of the recombinant rotavirus protein intramuscularly with the adjuvant QS-21 appeared to produce a greater IgG response than when the protein was administered intranasally in combination with the adjuvant LT. In contrast, intranasal administration with LT as the adjuvant produced a greater serum IgA response. Stool IgA was also greater when vaccination was performed intranasally using LT as the adjuvant.

#### Example 7

##### EIA Method to Measure Rotavirus Antigen in Stool to Determine Shedding

Stool specimens were thawed, homogenized and centrifuged (500 *g*, 10 min). For rotavirus antigen determination, 96-well EIA plates (Corning Costar Co.) were coated overnight at 4°C with 100 µl/ well of either rabbit antibody to rotavirus (duplicate positive wells) or preimmune rabbit serum (duplicate negative wells). Plates were washed and 50 µl of stool supernatant or serial two-fold dilutions of purified preparation of double-layered EDIM particles (standard) was added to duplicate wells coated with each antibody. After incubation at 37°C for one hour on

a rotation platform, plates were washed and 50  $\mu$ l normal goat serum (Vector) diluted 100-fold in phosphate-buffered saline containing 5% nonfat dry milk (PBS-M) was added for 15 minutes at room temperature. Fifty microliters of guinea pig antibody to rotavirus diluted 1:500 in PBS-M containing a 1:50 dilution of normal rabbit serum (DAKO) was added and incubated for 30 minutes. Plates were washed and 50  $\mu$ l of a 1:200 dilution of biotinylated goat anti-guinea pig IgG (Vector) in PBS-M containing a 1:50 dilution of normal rabbit serum was added and incubated 30 minutes. After washing plates, 50  $\mu$ l of a 1:100 dilution of peroxidase-conjugated avidin-biotin (Vector) in wash buffer was added and incubated 30 minutes. The plates were washed and 50  $\mu$ l substrate (*o*-phenylenediamine with H<sub>2</sub>O<sub>2</sub> in citric acid-phosphate buffer) was added and incubated (room temperature) for 15 minutes. The reaction was stopped with 75  $\mu$ l of 1.0 M H<sub>2</sub>SO<sub>4</sub>. The absorbance at 490 nm was measured and the net optical densities were determined by subtracting the average of the negative wells from the average of the positive wells. Values obtained from a standard curve generated from the serially diluted double-layered particles were used to determine concentrations of rotavirus protein in each specimen. The limit of detection was 3 ng/ml.

Reduction of shed rotavirus antigen was 79, 92 and almost 100% for MBP::VP6<sub>AB</sub>, MBP::VP6<sub>BC</sub> and MBP::VP6<sub>CD</sub>, respectively (Table 1). Therefore, the region CD was sufficient to elicit the same level of protection as the entire VP6 protein. It should be noted that MBP::VP6<sub>BC</sub> also provided excellent protection.

Table 1. Rotavirus antigen (ng/ml) shed in BALB/c mice immunized with MBP::VP6, MBP::VP6<sub>AB</sub>, MBP::VP6<sub>BC</sub> or MBP::VP6<sub>CD</sub>

Vaccine	Mouse Number	Day							Mean shedding per mouse per day	Reduction in shedding relative to control
		1	2	3	4	5	6	7		
None	1	0	25	122	131	88	395	13	42.66	
	2	0	0	0	0	0	0	0		
	3	0	0	48	569	49	7	23		
	4	0	0	14	22	23	25	0		
	5	0	0	19	60	222	23	20		
	6	0	0	19	85	19	9	0		
	7	0	0	25	161	25	8	7		
	8	0	0	17	58	28	24	6		
MBP::VP6	1	0	0	0	0	0	0	0	0.96	97.75%
	2	0	19	0	0	0	0	0		
	3	0	0	0	0	0	0	0		
	4	0	0	12	0	0	0	0		
	5	0	15	0	0	0	0	0		
	6	0	8	0	0	0	0	0		
	7	0	0	0	0	0	0	0		
	8	0	0	0	0	0	0	0		
MBP::VP6 <sub>AB</sub>	1	0	29	30	9	8	0	0	8.76	79.47%
	3	0	10	43	10	0	0	0		
	4	0	14	26	0	0	0	0		
	5	0	7	11	17	10	14	0		
	6	0	0	81	15	7	8	0		



Table 1. Rotavirus antigen (ng/ml) shed in BALB/c mice immunized with MBP::VP6, MPB::VP6<sub>AB</sub>, MBP::VP6<sub>BC</sub> or MBP::VP6<sub>CD</sub>

	Day									
	7	0	0	5	16	11	16	0		
	8	0	12	20	0	0	0	0		
MBP::VP6 <sub>BC</sub>	1	0	0	0	0	0	0	0	3.23	92.43%
	2	0	0	0	0	0	0	0		
	3	0	0	0	0	6	17	0		
	4	0	0	6	0	0	0	0		
	5	0	9	4	13	9	28	11		
	6	0	0	4	11	0	9	0		
	7	0	0	19	6	0	21	0		
	8	0	0	0	0	8	0	0		
MBP::VP6 <sub>CD</sub>	1	0	0	0	0	0	0	0	0.13	99.70%
	2	0	0	0	0	0	0	0		
	3	0	0	0	0	0	0	0		
	4	0	0	0	0	0	0	0		
	5	0	0	0	0	0	0	0		
	6	0	0	0	0	0	0	0		
	7	0	0	0	0	0	0	0		
	8	0	7	0	0	0	0	0		

A value of 0 indicates that shedding was below the limit of detection, i.e., < 3 ng/ml

#### Example 8

##### Comparison of the Adjuvants LT and CT

5 *E. coli* LT toxin was shown to be a very efficient adjuvant in inducing protection when inoculated intranasally with the rotavirus subunit vaccine MBP::VP6. (See Examples 4, 5, and 6). The effectiveness of *V. cholerae* toxin (CT) as an intranasal adjuvant was compared with LT. Mice were immunized with three doses of MBP::VP6 together with either 10 µg of CT or LT. Shedding data following virus challenge indicated that shedding was reduced by 98% irrespective of whether CT or LT was used (Table 2). Measurement of VP6-specific antibodies (Table 3)

10 revealed that specific IgG antibodies were induced by MBP::VP6 when co-administered with CT (GMT = 158,115 U/ml) or LT (GMT = 417,604 U/ml). Specific serum IgA was induced in CT- (GMT = 431 U/ml) and LT- (GMT = 1,185 U/ml) inoculated mice. Low but detectable stool IgA titers were also induced by CT (GMT = 41 U/ml) as well as LT (GMT = 77 U/ml). Therefore, both adjuvants could induce VP6 specific serum and mucosal antibodies.

Table 2. Rotavirus antigen shed (ng/ml) in BALB/c mice immunized with 3 doses of MBP::VP6 with *E. coli* LT or *V. cholerae* CT adjuvant

Adjuvant	Mouse Number	Day							Mean shedding per mouse per day	Reduction in shedding relative to control
		1	2	3	4	5	6	7		
None	1	0	25	122	131	88	395	13	42.66	
	2	0	0	0	0	0	0	0		
	3	0	0	48	569	49	7	23		
	4	0	0	14	22	23	25	0		
	5	0	0	19	60	222	23	20		
	6	0	0	19	85	19	9	0		
	7	0	0	25	161	25	8	7		
	8	0	0	17	58	28	24	6		
LT	1	0	0	0	0	0	0	0	0.96	97.75%
	2	0	19	0	0	0	0	0		
	3	0	0	0	0	0	0	0		
	4	0	0	12	0	0	0	0		
	5	0	15	0	0	0	0	0		
	6	0	8	0	0	0	0	0		
	7	0	0	0	0	0	0	0		
	8	0	0	0	0	0	0	0		
CT	1	0	0	0	0	0	0	0	0.71	98.34%
	2	0	0	0	0	0	0	0		
	3	0	0	0	0	0	0	0		
	4	0	0	0	0	0	0	0		
	5	0	0	0	0	0	0	0		
	6	0	0	0	0	0	0	0		
	7	0	0	0	0	0	0	0		
	8	0	18	0	0	13	0	0		

A value of 0 indicates that shedding was below the limit of detection, i.e., 3 ng/ml

Table 3. Rotavirus-specific antibodies (U/ml) in sera collected from BALB/c mice immunized with MBP::VP6 and with either LT or CT

Adjuvant	Mouse	Serum IgG		Serum IgA		Stool IgA	
		Pre-vaccination	Post-challenge	Pre-vaccination	Post-challenge	Pre-vaccination	Post-challenge
None	1	<100	9,710	<100	<100	<5	7,284
	2	<100	<100	<100	<100	<5	<100
	3	<100	5,674	<100	<100	<5	8,790
	4	<100	6,324	<100	<100	<5	4,885
	5	<100	4,233	<100	<100	<5	9,677
	6	<100	1,545	<100	<100	<5	2,511
	7	<100	4,596	<100	<100	<5	8,117
	8	<100	6,216	<100	<100	<5	2,567
LT	GMT		2,997				5,531
	1	<100	1,292,000	<100	995	<5	32,544
	2	<100	1,627,000	<100	2,401	<5	3,327
	3	<100	1,187,000	<100	560	<5	35,774
	4	<100	1,072,000	<100	720	<5	17,524
	5	<100	1,203,000	<100	816	<5	36,883
	6	<100	498,618	<100	1,373	<5	14,169
	7	<100	1,742,000	<100	1,035	<5	5,980
CT	8	<100	1,195,000	<100	3,486	<5	28,791
	GMT		1,162,701		1,185		16,720
	1	<100	226,207	<100	561	<5	4,863
	2	<100	213,608	<100	1,314	<5	10,501
	3	<100	202,023	<100	496	<5	4,178
	4	<100	178,661	<100	366	<5	21,325
	5	<100	69,919	<100	112	<5	4,634
	6	<100	225,320	<100	946	<5	24,361
	7	<100	220,965	<100	209	<5	6,575
	8	<100	64,345	<100	402	<5	42,188
	GMT		158,115		431		10,452
			934,476				41
							5,667
							6,500
							3,717
							5,260
							2,832
							5,431
							4,356
							4,953
							17,237
							9,515

Titers of &lt;100 U/ml indicate that no serum rotavirus IgG or IgA was detected.

Titers of &lt;5 U/ml indicate that no stool IgA was detected.

GMT = geometric mean titer

Example 9Optimization of the Immunization Protocol: Determination of the Number of Doses to Elicit a Protective Immunity

As seen in the previous examples, 2 or 3 doses of the MBP::VP6 induced almost complete protection.

To determine if 1 or 2 doses could provide the same level of protection, mice were immunized intranasally with 1, 2 or 3 doses of MBP::VP6 (8.8 µg/dose) using LT as adjuvant. For the latter two groups, doses were given at 14 days apart. Measurement of serum rotavirus-specific IgG indicated that the levels of IgG induced by three doses (GMT=417,604 U/ml) was higher than two doses (GMT= 122,839 U/ml), which in turn was higher than one dose (GMT=32,843 U/ml; Table 4). Serum IgA titers for 3 doses were higher (GMA= 1,185 U/ml) than 2 doses (GMT=256 U/ml) or 1 dose (GMT=243 U/ml). Larger titers of stool IgA could be detected in mice receiving 3 doses (GMT=77 U/ml) than 2 doses (GMT=24 U/ml). Only a few animals receiving 1 dose developed measurable stool rotavirus IgA (GMT= 12 U/ml).

Although the immunological responses differed between the 1, 2 and 3 dose protocols, animals were shown to be protected by a single vaccination. Analyses of the quantities of rotavirus antigen shed following rotavirus challenge one month after the last or only immunization indicated that 1, 2 or 3 doses of the vaccine resulted in almost 100, 98 and 98% reduction in shedding, respectively (Table 5). Therefore, one dose of MBP::VP6 was sufficient to induce essentially complete protection and protection appeared to be independent of the titer of specific antibodies.

Table 4. Rotavirus-specific antibodies in sera collected from mice immunized with 1, 2, or 3 doses of MBP::VP6

Adjuvant	Mouse Number	Serum IgG			Serum IgA			Stool IgA		
		Pre-vaccination	Pre-challenge	Post-challenge	Pre-vaccination	Pre-challenge	Post-challenge	Pre-vaccination	Pre-challenge	Post-challenge
None	1	< 100	< 100	9,710	< 100	< 100	7,284	< 5	17	4,517
	2	< 100	< 100	< 100	< 100	< 100	< 100	< 5	12	10
	3	< 100	< 100	5,674	< 100	< 100	8,790	< 5	13	5,920
	4	< 100	< 100	6,324	< 100	< 100	4,885	< 5	< 5	1,986
	5	< 100	< 100	4,233	< 100	< 100	9,677	< 5	16	2,325
	6	< 100	< 100	1,545	< 100	< 100	2,511	< 5	20	4,513
	7	< 100	< 100	4,596	< 100	< 100	8,117	< 5	8	3,576
	8	< 100	< 100	6,216	< 100	< 100	2,567	< 5	13	1,541
GMT				2,997			5,531		12	1,534
1 dose	1	< 100	64,219	804,286	< 100	468	90,536	< 5	39	10,982
	3	< 100	19,511	836,468	< 100	682	23,666	< 5	10	7,445
	3	< 100	29,575	53,079	< 100	188	415	< 5	27	33
	5	< 100	43,111	446,253	< 100	391	22,207	< 5	11	9,649
	6	< 100	30,789	442,076	< 100	166	20,193	< 5	6	5,108
	7	< 100	46,633	638,240	< 100	129	23,968	< 5	9	9,638
	8	< 100	17,970	908,344	< 100	< 100	37,835	< 5	6	7,318
GMT			32,843	455,751		243	16,696		12	3,694
2 doses	1	< 100	80,175	890,661	< 100	< 100	29,583	< 5	18	7,633
	2	< 100	80,966	731,137	< 100	223	80,418	< 5	13	7,096
	3	< 100	137,609	628,817	< 100	406	39,014	< 5	76	6,886
	4	< 100	194,994	1,887,000	< 100	578	98,965	< 5	93	13,548
	5	< 100	131,460	930,027	< 100	329	45,098	< 5	8	2,973
	6	< 100	164,938	865,635	< 100	131	43,500	< 5	16	6,599
	7	< 100	103,431	836,212	< 100	180	228,623	< 5	15	9,840
	8	< 100	132,712	1,148,000	< 100	456	68,520	< 5	35	10,097
GMT			122,839	937,588		256	64,023		24	7,485
3 doses	1	< 100	378,240	1,292,000	< 100	995	32,544	< 5	73	7,732
	2	< 100	311,525	1,627,000	< 100	2,401	3,327	< 5	57	4,717
	3	< 100	297,229	1,187,000	< 100	560	35,774	< 5	22	10,082
	4	< 100	273,683	1,072,000	< 100	720	17,524	< 5	30	4,550
	5	< 100	599,120	1,203,000	< 100	815	36,883	< 5	40	8,829
	6	< 100	481,863	498,618	< 100	1,373	14,169	< 5	658	4,690

Table 4. Rotavirus-specific antibodies in sera collected from mice immunized with 1, 2, or 3 doses of MBP::VP6

Adjuvant	Mouse Number	Serum IgG			Serum IgA			Stool IgA		
		Pre-vaccination	Pre-challenge	Post-challenge	Pre-vaccination	Pre-challenge	Post-challenge	Pre-vaccination	Pre-challenge	Post-challenge
	7	< 100	359,574	1,742,000	< 100	1,035	5,980	< 5	189	7,972
	8	< 100	929,583	1,195,000	< 100	3,486	28,791	< 5	90	5,771
	GMT		417,604	1,162,701		1,185	16,720		77	6,500

Titers of < 100 U/ml indicate that no serum rotavirus IgG or IgA was detected.

Titers of < 5 U/ml indicate that no stool IgA was detected

GMT=geometric mean titer

Table 5. Rotavirus antigen (ng/ml) shed in BALB/c mice immunized with EDIM after immunization with 1, 2 or 3 doses of MBP::VP

Number of Doses	Mouse Number	Day							Mean shedding per mouse per day	Reduction in shedding relative to control
		1	2	3	4	5	6	7		
None	1	0	25	122	131	88	395	13	42.66	
	2	0	0	0	0	0	0	0		
	3	0	0	48	569	49	7	23		
	4	0	0	14	22	23	25	0		
	5	0	0	19	60	222	23	20		
	6	0	0	19	85	19	9	0		
	7	0	0	25	161	25	8	7		
	8	0	0	17	58	28	24	6		
One dose	1	0	0	0	0	0	0	0	0.20	99.53%
	2	0	0	10	0	0	0	0		
	3	0	0	0	0	0	0	0		
	4	0	0	0	0	0	0	0		
	5	0	0	0	0	0	0	0		
	6	0	0	0	0	0	0	0		
	7	0	0	0	0	0	0	0		
	8	0	0	0	0	0	0	0		
Two doses	1	0	25	0	0	0	0	0	1.02	97.61%
	2	0	0	0	12	12	0	0		
	3	0	0	0	0	0	0	0		
	4	0	0	0	0	0	0	0		
	5	0	0	0	0	0	0	0		
	6	0	8	0	0	0	0	0		
	7	0	0	0	0	0	0	0		
	8	0	0	0	0	0	0	0		
Three doses	1	0	0	0	0	0	0	0	0.96	97.75%
	2	0	19	0	0	0	0	0		
	3	0	0	0	0	0	0	0		
	4	0	0	12	0	0	0	0		
	5	0	15	0	0	0	0	0		
	6	0	8	0	0	0	0	0		
	7	0	0	0	0	0	0	0		
	8	0	0	0	0	0	0	0		

A value of 0 indicates that shedding was below the limit of detection, i.e. < 3 ng/ml)

### Development of a Subunit Vaccine

To further facilitate the development of a safe and effective rotavirus vaccine candidate, the following work was performed to generate rotavirus fusion proteins containing an immunologically active fragment of VP6 for use as a vaccine candidate. Subunit vaccines are thought to be generally safer than killed virus or live-attenuated virus vaccines since only a portion of the virus is used to induce an immune response in a subunit vaccine, as opposed to the entire virus in either of the alternative methods. While use of entire viral proteins represents an advancement over whole virus vaccines, this approach could also be improved using partial protein or subunit vaccines. Such vaccines would reduce the cost of preparation and decrease the difficulties in preparing the needed quantities of viral protein to be used in the vaccine preparations. In light of this fact, the development of a bacteria-expressed subunit vaccine composed of a subset of those viral protein amino acids derivitized from these subunit vaccines that induce an immune response from an immunized individual would be advantageous. Accordingly, the examples below illustrate the methods used to generate a rotavirus fusion protein containing a subset of viral protein amino acids required to elicit an immune response from a vaccinated individual fused to a suitable fusion protein partner.

#### Example 10

#### Construction, Expression and Purification of Truncated VP6 Fusion Proteins

To produce a minimal subunit vaccine while retaining the original protective efficacy, three plasmids, pMAL-c2/EDIM6<sub>AB</sub>, pMAL-c2/EDIM6<sub>BC</sub> and pMAL-c2/EDIM6<sub>CD</sub>, were constructed to express truncated forms of VP6, wherein the truncated forms of VP6 contain immunogenic fragments of a rotavirus protein. Recombinant plasmids pMAL-c2/EDIM6<sub>AB</sub>, pMAL-c2/EDIM6<sub>BC</sub> and pMAL-c2/EDIM6<sub>CD</sub>, containing truncated forms of VP6 were constructed using the same strategy that was used for the construction of pMAL-c2/EDIM6, as seen in Example 1. These plasmids expressed MBP::VP6<sub>AB</sub> containing amino acids 1 to 196, MBP::VP6<sub>BC</sub> containing amino acid 97 to 297 and MBP::VP6<sub>CD</sub> containing amino acids 197 to 397. (See FIGURE 5). To construct these plasmids, cDNAs were synthesized by polymerase chain reaction (PCR) using pMAL-c2/EDIM6 (see Example 1) as the template. The gene specific primers used for construction and the regions of VP6 cloned are summarized in Table 6.

Table 6. Primers used to clone pMAL-c2/MBP<sub>AB</sub>, pMAL-c2/EDIM6<sub>BC</sub> and pMAL-c2/EDIM6<sub>CD</sub>

Plasmid	Name of Fusion Protein	Primers
pMAL-c2/MBP <sub>AB</sub>	MBP::VP6 <sub>AB</sub>	Forward primer: atg gat gtg ctg tac tct atc SEQ ID NO. 1 Reverse primer: tca cga gta gtc gaa tcc tgc aac SEQ ID NO. 2
pMAL-c2/EDIM6 <sub>BC</sub>	MBP::VP6 <sub>BC</sub>	Forward primer: atg gat gaa atg atg cga gag tca SEQ ID NO. 3 Reverse primer: tca gaa tgg cgg tct cat caa ttg SEQ ID NO. 4
pMAL-c2/EDIM6 <sub>CD</sub>	MBP::VP6 <sub>CD</sub>	Forward primer: tgc gca att aat gct cca gct SEQ ID NO. 5 Reverse primer: tca ctt tac cag cat gct tct aat SEQ ID NO. 6



Once constructed, the plasmids encoding the truncated VP6 fragments were introduced into bacteria for protein expression. Recombinant bacteria containing pMAL-c2/EDIM6<sub>AB</sub>, pMAL-c2/EDIM6<sub>BC</sub> and pMAL-c2/EDIM6<sub>CD</sub> were grown as described in the examples above. Specifically, an overnight culture was grown (37°C; shaken at 215 rpm) in rich broth (tryptone, 10 gm; yeast extract, 5 gm; NaCl, 5 gm; glucose, 2 gm; and ampicillin, 100 mg per liter). On the following day, 10 ml of overnight culture for each vector were inoculated into 1 liter of rich broth. The culture was grown until the optical density reached ~0.6 OD<sub>600</sub>. IPTG was added (0.3 mM) to induce expression of fusion protein. Growth was continued for 3 hours.

As previously described for MBP::VP6, cells expressing MBP::VP6<sub>AB</sub>, MBP::VP6<sub>BC</sub> or MBP::VP6<sub>CD</sub> were harvested by centrifugation (4,000 *g*; 20 minutes). The cells were washed in PBS and then recovered by centrifugation. The cell pellet was frozen at -20°C, thawed slowly in cold water, and resuspended in a total of 50 ml of buffer L (5 mM NaH<sub>2</sub>PO<sub>4</sub>, 10 mM Na<sub>2</sub>HPO<sub>4</sub>, 30 mM NaCl, 10 mM β-mercaptoethanol, 0.2% Tween 20, 1 mM PMSF, 25 mM benzamidine, and 200 μg/ml of lysozyme). After digestion for 15 minutes (room temperature), the suspension was sonicated by three 30 second bursts (BioSonic IV, 50% power setting) while placed in an ice/water bath. NaCl (265 mg) and RNase A (5 μl of 10 mg/ml) were added to each 10 ml of sonicated cell lysate. The lysate was then centrifuged (54,000 *g*, 30 minutes) to obtain a supernatant containing a crude preparation of fusion protein. Fusion proteins in the various crude preparation were purified by affinity chromatography as described for MBP::VP6. The individual supernatants containing the various fusion protein constructs were mixed with amylose resin (New England Biolabs, Beverly MA) for 2 hours in a 500-ml flask on a rocker. After centrifugation (2,100 *g*, 5 minutes), the resin was recovered, then resuspended in 50 ml of buffer C (Buffer L containing 0.5 M NaCl), rocked for 30 minutes and finally centrifuged to recover the resin. The resin was washed in this manner 3 times and then washed overnight with 500 ml of buffer C. On the following day, the resin was recovered by centrifugation and resuspended in 50 ml of buffer D (50 mM Tris-HCl, pH 7.5; 50 mM NaCl; 1 mM EDTA; 10 mM β-mercaptoethanol; 1 mM PMSF), and rocked for 30 minutes. The resin was spun down and the bound fusion protein was eluted from the resin by suspending the resin in 250 ml of 15 mM maltose in buffer D for 2 hours. The resin was recovered by centrifugation and the supernatant containing the fusion proteins was subjected to buffer exchange to PBS and was simultaneously concentrated by ultrafiltration using a stirred-cell concentrator (Amicon, Beverly MA; model 8400).

#### Example 11

##### Vaccination and Challenge of Mice Using Truncated VP6 Fusion Proteins

Six-week-old immunologically naïve female BALB/c mice (Harlan Sprague) and B cell deficient μMt mice were used to study the ability of the various truncated VP6 fusion proteins to elicit a protective response from vaccinated mice. Blood and stool specimens were collected from the animals prior to vaccination. Animals were immunized intranasally with 8.8 μg of fusion protein vaccines (MBP::VP6, MBP::VP6<sub>AB</sub>, MBP::VP6<sub>BC</sub> or MBP::VP6<sub>CD</sub>) in a 50-μl

volume. Animals, which received three doses, were immunized at biweekly intervals. Animals received 10 µg of the adjuvant LT with the vaccines. *E. coli* LT(R192G) was supplied by Dr. John Clements of Tulane University.

Four weeks after the last immunization, animals were bled and stool specimens were collected to measure antibody responses. Each animal was challenged with a 100- ID<sub>50</sub> dose, which is equivalent to 4 X 10<sup>4</sup> ffu, of EDIM virus (Lot number: P9 12/15/97). Two stool pellets were collected in 0.5 ml of Earl's Balanced Salt Solution (EBSS) from each mouse for seven days and stored at -20°C. Rotavirus antigen was measured in the stools by EIA to determine shedding. Twenty one days after challenge, sera and stool specimens were obtained again to measure antibody responses.

#### Example 12

##### Measurement of Serum Rotavirus IgG and IgA and Stool Rotavirus IgA

Serum rotavirus IgA and IgG and rotavirus stool IgA were measured by EIA. EIA plates (Corning Costar Co.) were coated overnight at 4°C with anti-rotavirus rabbit IgG. After washing with phosphate buffered saline plus 0.05% Tween 20, 50 µl of EDIM viral lysate or mock-infected cell lysate were each added to duplicate positive and duplicate negative wells for one hour at 37°C on a rotation platform. After washing plates, 50 µl of serial two-fold dilutions of pooled sera from EDIM infected mice assigned concentrations of 160,000 or 10,000 units/ml of rotavirus IgG or IgA, respectively, were added to duplicate wells coated with either EDIM-infected or uninfected MA104 cell lysates to generate a standard curve. Serial 10-fold dilutions of mouse sera to be tested were also added to duplicate wells of each lysate and incubated for 1 hour. This was followed by sequential addition of biotin-conjugated goat anti-mouse IgG or IgA (Sigma Chemical Co.), peroxidase-conjugated avidin-biotin (Vector Laboratories), and *o*-phenylenediamine substrate (Sigma Chemical Co.). Color development was stopped after fifteen minutes with 1 M H<sub>2</sub>SO<sub>4</sub> and the A<sub>490</sub> was measured. Titers of rotavirus IgG or IgA, expressed as units/ml, were determined from a standard curve generated by subtraction of the average A<sub>490</sub> values of the duplicate cell lysate wells from the average of the wells coated with EDIM lysate. For determination of stool rotavirus IgA, two stool pellets were collected into 0.5 ml of EBSS, homogenized, and centrifuged (1500 *g*, 5 min). Stool rotavirus IgA was then measured by the method described above.

Measurement of prechallenge VP6-specific antibody titers (Table 7) revealed that the AB and CD fragments induced high titers of serum IgG (GMT=43,625 U/ml and 129,920 U/ml respectively). Fragments AB and CD also induced serum IgA (GMT=222 U/ml and 716 U/ml respectively). Unexpectedly, no specific serum IgG or serum IgA (titer < 100) could be detected in MBP::VP6<sub>BC</sub>-inoculated mice by EIA (Table 7). Interestingly, the CD fragment induced specific stool IgA (GMT=70 U/ml) which was similar to the titer induced by the entire VP6 (GMT=77 U/ml, Table 3). Fragment AB induced marginally detectable titers while (GMT=20 U/ml) BC did not induce any stool IgA. Therefore, the protection generated with inoculation of BC appeared to be unrelated to stool rotavirus IgA.

Table 7. Rotavirus-specific antibodies in sera collected from BALB/c mice immunized with MBP:PVP6<sub>AB</sub>, MBP::VP6<sub>BC</sub> and MBP::VP6<sub>CD</sub> and LT

Adjuvant	Mouse number	Serum IgG			Serum IgA			Stool IgA		
		Pre-vaccination	Pre-challenge	Post-challenge	Pre-vaccination	Pre-challenge	Post-challenge	Pre-vaccination	Pre-challenge	Post-challenge
None	1	< 100	< 100	9,710	< 100	< 100	7,284	< 5	17	4,517
	2	< 100	< 100	< 100	< 100	< 100	< 100	< 5	12	10
	3	< 100	< 100	5,674	< 100	< 100	8,790	< 5	13	5,920
	4	< 100	< 100	6,324	< 100	< 100	4,885	< 5	< 5	1,986
	5	< 100	< 100	4,233	< 100	< 100	9,677	< 5	16	2,325
	6	< 100	< 100	1,545	< 100	< 100	2,511	< 5	20	4,513
	7	< 100	< 100	4,596	< 100	< 100	8,117	< 5	8	3,576
	8	< 100	< 100	6,216	< 100	< 100	2,567	< 5	13	1,541
GMT				2,997			5,531		12	1,534
MBP::VP6 <sub>AB</sub>	1	< 100	31,382	136,945	< 100	164	9,363	< 5	58	4,666
	3	< 100	37,245	65,499	< 100	260	11,332	< 5	40	5,381
	4	< 100	37,731	135,539	< 100	176	7,566	< 5	< 5	8,163
	5	< 100	43,943	134,255	< 100	195	9,970	< 5	6	7,401
	6	< 100	28,668	49,032	< 100	263	12,425	< 5	19	3,199
	7	< 100	63,053	260,311	< 100	151	14,237	< 5	23	9,513
	8	< 100	86,215	235,759	< 100	453	13,387	< 5	37	3,164
GMT			43,652	125,529		222	10,956		20	5,467
MBP::VP6 <sub>BC</sub>	1	< 100	100	66,661	< 100	< 100	1,470	< 5	13	1,859
	2	< 100	100	109,867	< 100	< 100	4,327	< 5	< 5	3,872
	3	< 100	100	72,780	< 100	< 100	9,919	< 5	7	1,762
	4	< 100	100	58,668	< 100	< 100	3,498	< 5	6	2,926
	5	< 100	100	60,548	< 100	< 100	8,879	< 5	< 5	6,602
	6	< 100	100	83,420	< 100	< 100	4,054	< 5	< 5	4,608
	7	< 100	100	126,528	< 100	< 100	11,466	< 5	< 5	7,061
	8	< 100	100	52,172	< 100	< 100	3,884	< 5	7	3,171
GMT							4,938		6	3,551
MBP::VP6 <sub>CD</sub>	2	< 100	155,158	420,416	< 100	957	266,389	< 5	60	18,245
	3	< 100	137,463	701,792	< 100	551	218,600	< 5	85	16,452
	4	< 100	137,679	524,277	< 100	696	65,597	< 5	84	11,163
	5	< 100	149,834	598,649	< 100	727	16,722	< 5	83	3,504
	6	< 100	89,976	212,449	< 100	979	54,195	< 5	157	9,141
	7	< 100	179,669	354,013	< 100	730	67,505	< 5	100	16,652
	8	< 100	87,840	555,205	< 100	508	67,927	< 5	15	7,249
GMT			129,920	452,205		716	76,880		70	10,377

Numbers of < 100 u/ml indicate that no serum rotavirus IgG or IgA was detected.

### Example 13

#### Western Blot Analysis

Serum samples from mice immunized with vaccines were analyzed for rotavirus protein-specific antibodies by Western blot analyses. Cesium chloride gradient-purified rotavirus particles were subjected to SDS-polyacrylamide gel electrophoresis. Separated rotavirus proteins were blotted to a nitrocellulose sheet and cut into strips each of which contained 3 µg of rotavirus proteins. The strips were blocked with 5% skim milk in Tris-HCl buffer (TBS, 50 mM Tris-

HCl, pH 7.5, 0.9% NaCl). The strips were then incubated with antisera obtained from immunized mice. After washing with 0.1% Tween-20 in TBS, the strips were incubated with goat anti-mouse IgG conjugated to alkaline phosphatase (Life Technologies, Gaithersburg, MD). The strips were washed with TBS and then incubated with 4-chloro-3-indolylphosphate and nitroblue tetrazolium (Life Technologies, Gaithersburg, MD) to visualize bound antibodies.

Western blot analyses confirmed that no specific antibodies could be detected in BMP::VP6<sub>BC</sub>-immune sera (FIGURE 6).

#### Example 14

##### Protection against EDIM Shedding by MBP::VP6 in $\mu$ Mt Mice

B-cell deficient  $\mu$ MT mice were also vaccinated intranasally with two doses (8.8  $\mu$ g/dose) of MBP::VP6 with LT. As expected, no rotavirus IgG, IgA or IgM was detected in the sera of any of these mice during this study. Analyses of virus shedding indicated that the subunit vaccine was as protective in these mice as was found with immunologically normal BALB/c mice (Table 8). This finding suggested that the vaccine could induce protection by a mechanism that did not require rotavirus antibodies. The mechanism was therefore not antibody dependent.

Table 8. Rotavirus shed in B-cell deficient Mt mice challenged with EDIM after immunization with 2 doses of MBP::VP6

Vaccine	Mouse Number	Day								Mean shedding per mouse per day	Reduction in shedding relative to control
		2	3	4	5	6	7	8	9		
None	1	1	11	39	197	114	10	0	0	764.44	
	2	98	3277	3579	2256	2760	1111	173	0		
	3	19	152	564	573	526	84	16	0		
	4	10	946	3588	2819	2065	1051	25	5		
	5	32	1227	2762	3271	92	20	0	0		
	6	377	505	1434	398	492	14	0	0		
	1	10	0	0	0	0	0	0	0	2.40	
	2	3	0	0	0	0	0	0	0		
	3	0	0	0	0	0	0	0	0		
	4	0	0	0	0	0	0	0	0		
	5	3	0	0	0	0	0	0	0		
	6	38	23	17	9	7	0	0	0		

Note: Shedding values of 0 indicate that shedding was below the limit of detection at 3 ng/ml.

In Example 15, two approaches were used to locate and characterize the protective epitopes in VP6. One approach was to immunize with expressed chimeric MBP proteins containing fragments of VP6. The other approach was to immunize with synthetic peptides. The goal was to identify the minimal protective epitopes which may possibly be incorporated into a subunit vaccine.

#### Example 15

##### Construction and Testing of Recombinant pMAL-c2 Plasmids Expressing

Fragments of BMP::VP6<sub>CD</sub>Minimal Subunit Vaccine

Because the C-terminal 50% of VP6 CD region could induce the same level of protection as the entire VP6 protein, the protective domains of this 201-amino acid portion of VP6 were further mapped. Four overlapping regions of the C-terminal CD region of the VP6 gene were cloned into pMAL-c2 (FIGURE 7). The specific primers that were used for construction of these plasmids are summarized in Table 9. As described previously, cDNAs were inserted into the restriction site *Xmn* I of pMAL-c2, placing the inserted sequences downstream from the *E. coli* Mal E gene, which encodes maltose binding protein (MBP), resulting in expression of MBP fusion proteins. These plasmids expressed MBP::VP6<sub>CD1</sub> containing amino acids 197 to 263, MBP::VP6<sub>CD2</sub> containing amino acids 244 to 310, MBP::VP6<sub>CD3</sub> containing amino acids 291 to 351, and MBP::VP6<sub>CD4</sub> containing amino acids 332 to 397 (FIGURE 8).

Recombinant plasmids were transformed into *E. coli*. White colonies of bacteria containing recombinant plasmids on agar plates were then identified in the presence of IPTG and X-gal, and selected for further screening by PCR for gene identity and orientation. Recombinant bacteria were grown as described previously for pMAL-c2/EDIM6. An overnight culture was grown (37°C; shaken at 215 rpm) in rich broth (tryptone, 10 gm; yeast extract, 5 gm; NaCl, 5 gm; glucose, 2 gm; and ampicillin, 100 mg per liter). On the following day, 10ml of overnight culture were inoculated into 1 liter of rich broth. The culture was grown until the A<sub>600</sub> reached ~0.6. IPTG was added (0.3 mM) to induce expression of the fusion protein. Growth was then continued for 3 hours. Nucleotide sequencing was used to ultimately confirm the authenticity of the rotavirus gene sequences.

The 201-amino acid long CD region of VP6 could induce the same level of protection as the entire VP6 protein. To further map protective domains in this portion of VP6, four overlapping regions are cloned into pMAL-c2. The sub-regions are designated CD1, CD2, CD3 and CD4, and contain 67, 67, 61 and 66 amino acids, respectively (FIGURE 7 & Table 9). The fusion proteins MBP::VP6<sub>CD1</sub>, MBP::VP6<sub>CD2</sub>, MBP::VP6<sub>CD3</sub> and MBP::VP6<sub>CD4</sub> containing these sub-regions are purified as described in Example 2 and tested for their protective efficacies.

Vaccination was performed as described in Example 15, Fine Mapping of Protective Epitopes using Synthetic Peptides. The protective efficiencies of various peptides were 88, 85, 19, and 92% for CD1, CD2, CD3, and CD4, respectively. These results show that peptides CD1, CD2, and CD4 have utility as vaccine components.

Table 9. Primers used to clone pMAL-c2/EDIM6<sub>CD1</sub>, pMAL-c2/EDIM6<sub>CD2</sub>, pMAL-c2/EDIM6<sub>CD3</sub> and pMal-c2c2/EDIM6<sub>CD4</sub>

<u>Plasmid</u>	<u>Name of fusion protein</u>	<u>Primers</u>
pMAL-c2/EDIM6 <sub>CD1</sub>	MBP::VP6 <sub>CD1</sub>	Forward primer: atg gat gtg ctg tac tct atc SEQ. I.D. NO. 7 Reverse primer: tca gaa ctc aac ttc tac att att tgg SEQ. I.D. NO. 8
pMAL-c2/EDIM6 <sub>CD2</sub>	MBP::VP6 <sub>CD2</sub>	Forward primer: gca act aca tgg tac ttc aac cca SEQ. I.D. NO. 9 Reverse primer: tca att tgg gaa aag tgc agt cac tgc SEQ. I.D. NO. 10
pMAL-c2/EDIM6 <sub>CD3</sub>	MBP::VP6 <sub>CD3</sub>	Forward primer: tca ttt caa ttg atg aga ccg cca SEQ. I.D. NO. 11 Reverse primer: tca ttg tct gac tga cgt cac att ggc SEQ. I.D. NO. 12
pMAL-c2/EDIM6 <sub>CD4</sub>	MBP::VP6 <sub>CD4</sub>	Forward primer: gaa tca gtt ctc gcg gat gca agt SEQ. I.D. NO. 13 Reverse primer: tca ctt tac cag cat gct tct aat SEQ. I.D. NO. 14

#### Fine Mapping of Protective Epitopes using Synthetic Peptides

This approach was used to determine the smallest subunit vaccine possible. Synthetic peptides were designed to identify the protective domain(s) in the carboxyl-terminal half or CD region of VP6. A series of 11 overlapping peptides (Table 10) were synthesized by Quality Controlled Biochemicals, Inc (Hopkinton, MA). The synthetic peptides were on a Perkin Elmer 9050 peptide synthesizer or were synthesized manually. The well-established solid phase method was employed utilizing orthogonally protected amino acids. Cleavage and deprotection were done in aqueous trifluoroacetic acid. These overlapping peptides contained between 18 and 31 amino acids. All 11 peptides were tested as immunogens with LT(R192G).

Six-week-old rotavirus antibody-free female BALB/c mice (Harlan Sprague) and B-cell deficient Mt mice were used for vaccination. Blood and stool specimens were collected from the animals prior to vaccination. Animals were immunized intranasally with 8.8 g of fusion proteins or 50 g of synthetic peptides in a 50- $\mu$ l volume. Animals receiving either two or three doses were immunized at biweekly intervals. The adjuvant *E. coli* LT(R192G) (10 g supplied by Dr. Clements of Tulane University) was coadministered with the test vaccine.

Four weeks after the last immunization animals were bled and stool specimens were collected to measure antibody response. Each animal was challenged with a 100 ID<sub>50</sub> dose, which is equivalent to  $4 \times 10^4$  ffu of EDIM virus, passage 9. Two stool pellets were collected into 1.0 ml of Earle's balanced salt solution (EBSS) from each mouse for seven or more days and stored at -20°C. Rotavirus antigen was measured in the stool by EIA to determine shedding (See Example 7). Twenty-one days after challenge, sera and stool specimens were obtained again to measure antibody responses.

The protective efficacies of seven peptides (3, 5, 6, 7, 9, 10, and 11) were first examined in BALB/c mice. It was found that two immunizations [50 g with 10 g of LT(R192G)] with peptides 6, 11, or 3 induced a mean reduction in rotavirus shedding of 88, 64, and 57% respectively (Table 15,  $P < .001$ ). The other 4 peptides did not elicit protection. Peptide 6, a 25mer, contains a 14-amino acid sequence RLSFQLMRPPNMTP that has been identified by a proliferation assay to be an H-2<sup>d</sup> CD4 epitope (Banos, et al., *J. Virol.* 71:419-426, 1997). This 14mer peptide, tentatively called 6-14, was also synthesized. Experiments using this 14mer indicated that it provided comparable protection to that of peptide 6 in H-2<sup>d</sup> BALB/c mice.

The protective efficacies of peptide 1, 2, and 4 were tested. Peptides 2 and 4 were found to be 70 and 77% protective, respectively. The excellent protection observed with 6 of the peptides demonstrated that the maltose-binding protein is not required for induction of protection. These findings were confirmed with peptides 2 and 4. These results suggest the possibility of an alternative or supplemental vaccine strategy, i.e. a multi-peptide vaccine, which can be formulated from a selection of protective epitopes.

Subsequently, B-cell deficient  $\mu$ Mt (H-2<sup>b</sup>) mice are used to determine whether or not peptide(s)-induced antibodies are involved in protection.

Table 10. Synthetic peptides for mapping protective domains in the CD region of VP6

Peptide Number	Sequence	SEQ. I.D. NO.
#1	CAINAPANIQQFEHIVQLRRVLT	SEQ. I.D. No. 15
#2	PDAERFSFPRVINSADGA	SEQ. I.D. No. 16
#3	FSFPRVINSADGATTWYFNPVILRPNNVEV	SEQ. I.D. No. 17
#4	FNVPILRPNNVEVEFLNGQVINTYQARF	SEQ. I.D. No. 18
#5	NGQVINTYQARFGTIVARNFDITIRLSFQLM	SEQ. I.D. No. 19
#6	RNFDITIRLSFQLMRPPNMTPAVTAL	SEQ. I.D. No. 20
#7	MTPAVTALFPNAQPFHHATVGLTLRIDSA	SEQ. I.D. No. 21
#8	HATVLTLRIDSAICESVLADASETMLANV	SEQ. I.D. No. 22
#9	VLADASETMLANVTSVRQEYAI	SEQ. I.D. No. 23
#10	QEYAIQVGPVFPFGMNWTDLITNYSRPSRED	SEQ. I.D. No. 24
#11	TDLITNYSRPSREDNLQRVFTVASIRSMVLK	SEQ. I.D. No. 25

#### Example 16

##### Mapping of Antibody Independent Epitopes Using Mt mice

Understanding the mechanisms of protection is crucial to vaccine development. A unique observation obtained with chimeric vaccine, disclosed here, is that B-cell deficient Mt mice were found to be as well protected from shedding following vaccination with MBP::VP6 and LT(R192G) as immunologically normal BALB/c mice (See Example 14). This finding suggests that VP6 may induce completely antibody-independent protective immunity, a phenomenon that has not been reported previously for a rotavirus vaccine candidate. To locate the epitopes

responsible for this protective mechanism, an experiment was conducted to initially examine the AB and CD peptides (i.e., MBP fusion proteins containing the first and second halves of VP6, respectively). Because peptide 6 provided excellent protection (88% reduction in shedding) and contains a putative CD4 epitope, groups of Mt mice were also inoculated with peptide 6 and 6-14, both of which contain the putative CD4 epitope.

Examples 17 and 18 describe further experiments to determine which T cells are involved in immunity.

#### Example 17

##### The role of CD8 T cells in MBP::VP6-mediated immunity

Studies using rotavirus particles for intranasal immunization have shown that CD8 cells are not needed for protection. Experiments were performed to determine whether immunization with VP6 can also mediate CD8-independent protection. To achieve this goal, Mt mice were depleted of CD8 by injection with anti-CD8 antibodies. These mice were then immunized intranasally with MBP::VP6 and LT(R192G) and challenged with EDIM, as discussed above. The results showed that these animals were equally protected from viral infection following immunization whether or not CD8 cells were present or removed at the time of challenge.

#### Example 18

##### The role of CD4 cells in protection

Based on results found with Mt mice, protection stimulated by VP6 was found not to be dependent on antibody production. Furthermore, protection following immunization with MBP::VP6 was not dependent on CD8 cells. This would leave CD4 cells as the most likely memory cells involved in protection. The importance of CD4 cells in protection following i.n. immunization using monoclonal antibody depletion as well as using genetically altered mice that lack CD4 cells was undertaken using the methods discussed above. In preliminary studies, it was found that i.n. immunization with double-layered rotavirus particles is much less protective in CD4-depleted or CD4 knock-out mice than in non-depleted or genetically normal mice. Similar results were obtained with CD4-depleted BALB/c mice immunized i.n. with the 6-14 peptide. It should be noted that peptide 6-14 discussed above provided nearly complete protection and this 14 amino acid peptide contains a known CD4 epitope.

#### Example 19

##### Rotavirus Fusion Protein Containing A Polymeric Histamine Fusion Partner (pMAL-c2X/EDIM6-6his)

##### Construction of pMAL-c2X/EDIM6-6his

To facilitate purification of a full length chimeric VP6, the plasmid pMAL-c2X (New England Biolabs, Beverly, MA) was used. The same construct can be used to express MBP::VP6::6Xhis or VP6::6Xhis. pMAL-c2X contains an *Nde* I site just 5' to the *Mal* E gene. This restriction site is one of 2 sites needed to delete the *mal* E sequence for the construction of pMAL-c2X/EDIM6-His6 that expresses the fusion protein 6his::VP6. The forward primer contains an *Nde* I site and a 5' terminal sequence of VP6 (nucleotides 4 – 21). The reverse sequence contains a stop codon (taa), 6 histidine-encoding codons and a 15-nucleotide VP6 carboxyl terminus sequence (Table 11). The newly added C-terminal 6Xhis fusion tag together with the N-terminal MBP enable the purification of the full-length VP6 using consecutive amylose resin and Talon (Palo Alto, CA) resin affinity chromatography. To create pMAL-c2X/EDIM6-6his, the



recombinant plasmid was modified by digestion with *Nde* I and re-ligated to create the plasmid pc2X/EDIM6-6his. This plasmid expresses VP6::6Xhis that is devoid of MBP.

Purification of full length MBP::VP6::6Xhis

MBP::VP6::6Xhis was expressed as described above for the fusion peptides of CD and purified using amylose resin, also described (Example 15). It was found that the purified preparation also contained truncated MBP::VP6 lacking the C-terminal 6 histidine residues and varying lengths of the VP6 terminal. Full-length MBP::VP6::6Xhis was then purified from the truncated proteins using Talon affinity resin containing the divalent cobalt which selectively binds hexahistidines (Clontech, Palo Alto CA). To do this, the protein samples were denatured by adding guanidine-HCl, Tris-HCl (pH 8) and NaCl (final concentrations of 6M, 50 mM and 400 mM, respectively). Talon resin was washed twice with sample lysis buffer (6 M guanidine-HCl, 400 mM NaCl, 50 mM Tris-HCl, pH 8). The protein solution was added to the washed Talon resin. The mixture was then gently agitated for at least 2 hours on a platform shaker. The resin was spun down in a centrifuge (700 g, 5 min) and the supernatant was discarded. The resin was washed by adding 10 bed-volumes of lysis buffer to the resin and the mixture was again agitated for 10 min on a platform shaker. The resin was spun down as before (700 g, 5 min) and the supernatant was discarded. The resin was washed in this way for a total of 4 times. The resin was resuspended in 1 bed-volume of lysis buffer and transferred to a 2-ml gravity-flow column. The resin was then washed twice with a wash buffer containing 8 M urea, 400 mM NaCl, 50 mM Tris-HCl, pH 8. The bound MBP::VP6::6Xhis was then eluted from the resin with elution buffer (6 M urea, 100 mM NaCl, 200 mM imidazole, 500 mM EDTA, 50 mM Tris-HCl, pH 8). The eluted protein was subjected to buffer exchange to PBS by using Centrprep 50 filters (Amicon Inc., Beverly MA).

Western blot analysis of MBP::VP6::6Xhis. Purified MBP::VP6::6Xhis protein was analyzed by Western blot analysis to determine its purity. Purified MBP::VP6::6Xhis protein was subjected to SDS-polyacrylamide gel electrophoresis and blotted onto a nitrocellulose sheet. The sheet was blocked with 5% skim milk in Tris-HCl buffer (TVS; 50 mM Tris-HCl, pH 7.5, 0.9% NaCl). Duplicate sheets were then incubated with anti-MBP (New England Biolabs, Inc., Beverly MA) or anti-6Xhis (Santa Cruz, CA) sera. After washing with 0.1% Tween-20 in TBS, the strips were incubated with goat anti-rabbit IgG conjugated to alkaline phosphatase (Life Technologies, Gaithersburg, MD). The strips were washed with TBS and then incubated with 4-chloro-3-indolylphosphate and nitroblue tetrazolium (Life Technologies, Gaithersburg, MD) to visualize bound antibodies. Preliminary experiments revealed a single protein corresponding to MBP::VP6::6Xhis that appeared to free of major contamination by truncated proteins.

Immunization of mice with MBP::VP6::6Xhis. Groups of 8 BALB/c mice have been immunized with 2 sequential doses (8.8 mg/dose) of the purified, doubly tagged VP6 protein and challenged with EDIM 1 month after the last immunization. Results from these experiments showed that mice immunized with this fusion protein were protected from rotaviral disease to a degree comparable to that found with MBP::VP6 immunized mice.

Table 11. Primers used to clone pMAL-c2X/EDIM6-6his

Primer	Sequence
Forward	<u>cat atg</u> <sup>1</sup> gac gtg ctg tac tct atc SEQ. I.D. NO. 26
Reverse	tta <u>atg atg atg atg atg atg</u> <sup>2</sup> ctt tac cag cat gct SEQ. I.D. NO. 27

<sup>1</sup> *Nde*I restriction site is underlined<sup>2</sup> codons for His

To analyze whether MBP is absolutely necessary for immunization, a VP6::6Xhis fusion protein was constructed. VP6::6Xhis, although still a fusion protein is considerably less bulky. Example 20 presents the construction and analysis of this protein.

#### Example 20

##### Recombinant Rotavirus Fusion Protein Replacing MBP with 6his

The construction of a recombinant rotavirus fusion protein using a fusion partner of 6 histidines rather than MBP is described below. Although MBP by itself did not induce protection or rotavirus-specific antibodies, it is not clear if it can modulate VP6-induced protective immunity. To determine whether MBP has any adjuvant effects on the protective efficacy of VP6, as well as to show the ability of other amino acid sequences to serve as fusion protein partners, a recombinant plasmid was constructed by first cloning VP6 into pMAL-c2X, as described above. This plasmid is identical to pMAL-c2 except that it contains a *Nde*I restriction site which, together with an engineered *Nde*I site (Table 11), allows the eventual deletion of the *malE* sequence.

Characterization of the recombinant plasmid shows the authenticity of the coding region. The 6his::VP6 recombinant rotavirus fusion protein expresses well, compared to the expression levels of the other recombinant rotavirus fusion proteins. The presence of the 6his sequence may prevent the assembly of the recombinant rotavirus fusion protein into a multimeric form and facilitates the purification of the recombinant protein. The efficacy of 6his::VP6 to elicit a protective immune response from an individual immunized with a vaccine containing this protein is compared with that of MBP::VP6. The results of this comparison show that the two recombinant rotavirus fusion proteins are capable of eliciting protective immune responses.

Even though intranasal immunization with VP6 provided nearly complete protection against rotavirus shedding following a subsequent challenge with murine rotavirus, the duration of protection had not been examined. In Example 21 the duration of protection using the MBP::VP6 is presented.

Further experiments were performed to optimize vaccination using the present vaccines. Examples 21-24 present data analyzing duration, other mucosal routes, other adjuvants, and the effect of dosage on the immune response.

#### Example 21

##### Duration of Protection Experiments

In the typical immunization protocol, mice were challenged 1 month after the last immunization. For this study, the time between the last immunization and challenge was extended to 3 months to determine whether the degree of protection (quantity of virus shed after challenge) is reduced with time. Mice given two intranasal immunizations with MBP::VP6 (8.8 g/dose) and LT(R129G) separated by a 2 week interval were found to be equally protected at 3 months (99.7%) or 1 month (97.8%) after the immunization (Table 12). This finding demonstrated that protection is not rapidly lost after immunization, an important finding regarding the utility of VP6 as a vaccine candidate.

Table 12. Protection of BALB/c mice at 1 versus 3 months after i.n. immunization with 2 doses of MBP::VP6

Table 12: Protection of BALB/c mice at 1 versus 3 months after i.m. immunization with 2 doses of M21-LT-G										
	Mouse number	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7	Shedding per mouse per day	% Reduction in shedding
1 month										
Unimmunized	1	0	64	2,050	470	269	52	0	315	
	2	0	92	1,508	185	159	32	0		
	3	0	103	272	194	196	9	0		
	4	0	18	3,740	1,132	603	88	0		
	5	0	18	1,051	78	61	23	0		
	6	0	59	2,761	222	86	23	0		
	7	0	8	169	238	66	55	0		
	8	0	0	59	1,026	290	105	0		
VP6+LT (R192G)	1	0	15	19	0	0	0	0	7	97.8
	2	0	8	13	7	19	4	0		
	3	0	6	8	0	23	0	0		
	4	0	21	46	19	43	0	0		
	5	0	14	11	0	0	0	0		
	6	0	8	0	0	0	0	0		
	7	20	90	0	0	0	0	0		
	8	0	20	0	0	0	0	0		
3 months										
Unimmunized	1	0	5	987	621	310	80	8	592	0
	2	0	177	711	167	108	15	0		
	3	3	11	538	1,298	560	116	4		
	4	0	22	233	262	122	40	0		
	5	0	1,739	12,566	1,650	996	377	8		
	6	0	61	397	303	219	17	0		
	7	0	169	3,179	1,426	434	137	8		
	8	0	15	349	1,922	532	261	4		
VP6+LT (R192G)	1	0	15	0	0	0	0	0	2	99.7
	2	0	0	10	0	0	0	0		
	3	3	0	0	0	0	0	0		
	4	0	19	17	0	0	0	0		

Table 12. Protection of BALB/c mice at 1 versus 3 months after i.n. immunization with 2 doses of MBP::VP6

Mouse number	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7	Shedding per mouse per day	% Reduction in shedding
5	0	0	9	4	0	0	0		
6	0	0	0	0	0	0	0		
7	0	6	0	0	0	0	0		
8	0	9	0	0	0	0	0		

Example 22Induction of protective immunity by another mucosal route

To determine whether MBP::VP6 is protective if delivered by a mucosal route other than intranasally, groups of mice were immunized orally with 2 inoculations of MBP::VP6 (8.8 g per inoculation), either with or without LT(R192G). Another group was immunized intranasally with this fusion protein and LT(R192G) for comparison. Immunized mice were challenged with murine rotavirus 1 month after the last immunization and the percent reduction in viral shedding was calculated (Table 13). Oral immunization with MBP::VP6 and LT(R192G) induced good protection (85% reduction in shedding) but this reduction was significantly ( $P < .001$ ) less than after i.n. immunization (99%). Therefore, i.n. was more effective than oral immunization; however, it is possible that the two routes may be used concomitantly to increase protection, a possibility to be examined in future experimentation. Induction of protection by oral inoculation, as in the case of intranasal immunization, was dependent on the presence of LT(R192G), which reemphasized the requirement for an adjuvant to be used in conjunction with the VP6 vaccine.

Table 13. Protection of BALB/c mice by oral versus i.n. immunization with 2 doses of MBP::VP6

Immunogen	Mouse number	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7	Mean shedding per mouse per day	% Reduction in shedding
<b>Oral</b>										
Unimmunized	1	0	11	862	1,190	208	79	0	385	
	2	0	363	3,491	592	508	62	0		
	3	0	220	2,491	846	184	116	4		
	4	0	85	943	490	184	221	24		
	5	0	62	740	584	110	146	0		
	6	0	41	2,480	1,248	652	151	0		
	7	0	6	242	663	118	108	0		
	8	0	20	217	532	137	117	4		
VP6	1	0	381	4,579	412	308	44	0	571	0
	2	0	166	1,01	199	192	10	0		

Table 13. Protection of BALB/c mice by oral versus i.n. immunization with 2 doses of MBP::VP6

Immunogen	Mouse number	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7	Mean shedding per mouse per day	% Reduction in shedding
VP6+ LT(R192G)	3	5	893	1,531	260	184	6	0	59	85
	4	0	698	4,602	592	148	8	0		
	5	0	244	1,895	575	120	34	0		
	6	0	757	3,424	1,823	816	380	22		
	7	0	16	478	430	189	86	16		
	8	0	188	255	352	50	19	0		
	2	0	75	102	3	0	10	0		
	3	0	149	108	11	3	4	0		
	4	0	132	156	43	11	14	0		
	5	0	93	899	196	45	33	0		
	6	0	14	6	0	0	0	0		
	7	0	118	121	33	8	10	0		
	8	10	25	15	0	0	0	0		
Intranasal										
VP6+ LT(R192G)	1	0	5	5	3	20	0	0	5	99
	2	0	0	8	0	0	0	0		
	3	0	29	0	0	0	0	0		
	4	0	16	5	0	0	0	0		
	5	0	13	8	0	0	0	0		
	6	0	68	26	3	0	0	0		
	7	0	14	10	0	0	0	0		
	8	0	7	0	0	23	0	0		

Example 23Effect of Different Adjuvants on Protection

Six-week-old rotavirus antibody-free female BALB/c mice (Harlan Sprague) and B-cell deficient Mt mice were used for vaccination. Blood and stool specimens were collected from the animals prior to vaccination. Animals were immunized orally or intranasally with 8.8 g of fusion proteins. Animals, receiving either two or three doses were immunized at biweekly intervals. When adjuvants were coadministered with the test vaccine, *E. coli* LT(R192G) (10 g supplied by Dr. Clements of Tulane University), *V. cholerae* CT (10 mg Sigma Chemical Co., St. Louis, Mo), poly[di(carboxylatophen-oxy)phosphazene] (PCPP 50 g, Avant Immunotherapeutics, Needham, MA) or QS-21 (20 g Wyeth-Lederle Laboratories) was used.

Four weeks after the last immunization, animals were bled and stool specimens were collected to measure antibody response. Each animal was challenged with a 100 ID<sub>50</sub> dose, which is equivalent to 4 X 10<sup>4</sup> ffu, of EDIM

virus passage 9. Two stool pellets were collected into 1.0 ml of Earle's balanced salt solution (EBSS) from each mouse for seven or more days and stored at -20°C. Rotavirus antigen was measured in the stool by EIA to determine shedding (See Example 7). Twenty-one days after challenge, sera and stool specimens were obtained again to measure antibody responses.

It had already been shown that cholera toxin (CT), which is biologically and functionally related to LT, could replace LT(R192G) as adjuvant. The effectiveness of other adjuvants (PCPP, QS-21) have now been examined. Groups of mice were vaccinated with 2 i.n. immunizations (two weeks apart) with MBP::VP6 and adjuvant [PCPP, QS-21 or LT(R192G)] (Table 14). The adjuvant PCPP conferred an 80% reduction in shedding when administered intranasally with MBP::VP6 but it was not effective when given orally. In contrast, 59% and 43% reductions were observed when QS-21 was included with MBP::VP6 for oral and intranasal inoculation, respectively. As previously observed, LT(R192G) provided 99% protection when given with the vaccine intranasally but was less protective (85%) when administered orally. These results indicate that the choice of adjuvant and the route of mucosal inoculation both impact the efficacy of the VP6 vaccine.

To search for other effective adjuvants, nucleic acid adjuvants (CpG DNA from CpG ImmunoPharmaceuticals, Wellesley, MA), double-stranded RNA, and the cholera toxin A1-based gene fusion protein CTA1-DD (Agren, et al. J Immunol 162:2432-2440, 1999), will be tested for their effectiveness in stimulating VP6-induced protective immunity.

Table 14. Effect of different adjuvants on oral and intranasal MBP::VP6-induced protection of BALB/c mouse

Immunogen	Mouse number	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7	Shedding per mouse per day	% Reduction in shedding
<b>Intranasal</b>										
Unimmunized	1	0	208	835	70	104	37	0	311	
	2	0	37	674	249	623	117	10		
	3	0	88	2,838	685	1,686	588	31		
	4	0	193	552	574	242	64	0		
	1	0	20	420	540	247	207	5		
	2	0	28	2,901	481	159	167	10		
	3	0	19	962	198	134	59	13		
	4	0	0	0	30	176	133	12		
VP6+LT(R192G)	1	0	0	0	0	0	0	0	2	99
	2	0	10	0	0	0	0	0		
	3	0	13	0	0	0	0	0		
	4	0	7	0	0	0	0	0		
	5	7	8	0	0	0	0	0		
	6	0	29	0	0	0	0	0		
	7	0	14	5	0	0	0	0		
	8	0	9	7	0	0	7	0		
VP6+QS21	1	0	7	285	1,200	263	62	0	177	43
	2	0	47	570	200	365	19	0		
	3	0	6	159	206	291	34	9		
VP6+PCPP	1	0	24	86	43	43	0	0	63	80

Table 14. Effect of different adjuvants on oral and intranasal MBP::VP6-induced protection of BALB/c mouse

Immunogen	Mouse number	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7	Shedding per mouse per day	% Reduction in shedding
Oral	2	0	97	220	138	324	14	0	385	85
	3	0	15	203	100	15	0	0		
	4	0	51	260	21	97	0	0		
	5	5	18	339	54	69	19	0		
	6	6	189	76	29	42	0	0		
	7	0	30	20	30	23	0	0		
	8	14	200	193	221	250	0	0		
	8	14	200	193	221	250	0	0		
Unimmunized	1	0	11	862	1,190	208	79	0	385	85
	2	0	363	3,491	592	508	62	0		
	3	0	220	2,491	846	184	116	4		
	4	0	85	943	490	184	221	24		
	5	0	62	740	584	110	146	0		
	6	0	41	2,480	1,248	652	151	0		
	7	0	6	242	663	118	108	0		
	8	0	20	217	532	137	117	4		
VP6+LT (R192G)	1	0	188	255	352	50	19	0	59	85
	2	0	75	102	3	0	10	0		
	3	0	149	108	11	3	4	0		
	4	0	132	156	43	11	14	0		
	1	0	93	899	196	45	33	0		
	2	0	14	6	0	0	0	0		
	3	0	118	121	33	8	10	0		
	4	10	25	15	0	0	0	0		
VP6+QS-21	1	0	0	56	169	20	37	6	156	59
	2	0	131	175	35	0	4	0		
	3	0	25	505	801	115	47	0		
	4	0	42	562	714	506	303	12		
	1	0	0	546	269	55	14	0		
	2	0	159	1,066	208	66	20	0		
	3	0	36	627	378	34	39	0		
	4	0	14	153	204	536	48	11		
VP6+PCPP	1	0	77	221	271	80	22	0	387	0
	2	0	279	665	3,227	91	485	0		
	3	0	126	2,656	436	4,816	8	3		
	4	0	128	375	124	33	14	0		
	1	0	125	542	210	129	12	0		
	2	0	48	2,249	145	22	8	0		
	3	0	316	1,222	150	440	35	0		
	4	0	75	1,033	302	440	27	3		

Table 15. Induction of protection by i.n. vaccination with overlapping synthetic peptides of the CD region of VP6 with LT (R192G)

Peptides	Mouse number	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7	Shedding per mouse per day	% Reduction in shedding
Unimmunized	1	0	10	187	418	333	102	0	225	
	2	0	54	238	444	129	80	0		
	3	0	33	706	1,251	1,046	99	0		
	4	0	15	312	297	291	52	0		
MBP::VP6CD	1	0	79	767	265	240	40	0	3	99
	2	0	47	429	860	589	409	0		
	3	0	23	767	387	412	145	0		
	4	0	19	202	264	218	354	0		
	1	7	16	0	0	0	0	0		
	2	0	0	16	13	0	0	0		
	3	5	19	0	0	0	0	0		
	4	0	0	0	0	0	0	0		
	1	0	10	5	11	0	0	0		
	3	0	26	0	0	0	0	0		
	4	0	0	19	0	0	0	0		
	1	0	21	300	86	24	17	0		
Peptide 3	2	26	294	472	134	175	16	0	96	57
	3	15	144	207	148	37	17	0		
	4	5	140	292	37	52	14	0		
	1	10	129	362	598	395	23	0		
Peptide 5	2	10	6	196	71	20	18	0	341	0
	3	11	90	53	9	15	11	0		
	4	20	94	434	86	44	14	0		
	1	12	128	261	270	150	31	0		
	3	15	84	1,162	687	276	44	0		
	4	0	25	629	373	272	43	18		
	1	4	70	630	268	767	45	9		
	2	5	148	988	666	1,738	409	7		
Peptide 6	3	25	167	2,684	982	1,437	102	0	27	88
	4	7	36	480	263	234	58	0		
	1	12	76	23	6	26	34	0		
	2	0	40	71	27	24	43	0		
	3	6	25	29	0	11	12	0		
	4	0	97	38	9	23	24	0		
	1	11	29	52	7	10	13	0		
	2	23	22	22	45	39	13	0		
	3	6	25	29	0	11	12	0		
	4	0	97	38	9	23	24	0		
	1	11	29	52	7	10	13	0		
	2	23	22	22	45	39	13	0		
	3	0	58	40	4	24	25	0		
	4	0	21	264	76	35	20	0		



Table 15. Induction of protection by i.n. vaccination with overlapping synthetic peptides of the CD region of VP6 with LT (R192G)

Peptides	Mouse number	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7	Shedding per mouse per day	% Reduction in shedding
Peptide 7	1	24	110	279	49	88	25	0	185	18
	2	0	17	990	402	178	115	16		
	3	0	164	555	117	90	42	0		
	4	0	0	428	288	200	130	0		
	1	7	123	1,845	144	35	53	0		
	2	0	48	1,063	293	692	242	9		
	3	0	45	401	110	77	32	0		
	4	15	147	486	97	69	26	0		
Peptide 9	1	63	277	2,707	249	66	41	0	275	0
	2	0	235	2,131	1,024	510	53	8		
	3	0	163	1,714	289	111	37	0		
	4	0	38	275	118	31	12	0		
	1	0	76	393	270	205	49	0		
	2	20	148	939	371	149	14	0		
	3	0	68	948	364	209	39	0		
	4	0	152	385	265	158	34	0		
Peptide 10	1	0	134	906	177	77	30	0	1825	0
	2	0	22	998	1,921	1,06	186	0		
					7					
	3	0	85	1,428	287	256	53	0		
	4	0	13	226	296	317	62	0		
	1	8	479	59,67	14,71	6,85	1,261	0		
				3	8	1				
	2	0	110	985	1,379	669	96	0		
Peptide 11	3	8	222	991	5,318	65	36	0	80	64
	4	6	168	209	286	85	24	0		
	1	0	54	277	47	23	30	0		
	2	0	19	186	220	47	29	0		
	3	0	17	297	136	51	30	0		
	4	15	31	487	59	39	41	0		
	1	0	49	294	230	98	36	0		
	3	0	32	212	181	62	23	0		

Example 24Effect of Dosage on Protection

The effect of dosage (1.76 g and 8.8 g) on protection by i.n. immunization with MBP::VP6 and LT(R192G) was examined (Table 16). Although mice immunized with two 1.76 g dosages plus LT(R192G) of chimeric VP6 appeared to be nearly as well protected as those administered two 8.8 µg-doses, (protective levels were 94 and 99%, respectively) the 94% protection level was significantly ( $P = .0003$ ) lower than the 99% protection level.

Table 16. Effect of 2 i.n. immunizations of 1.76  $\mu$ g or 8.8- $\mu$ g dose of MBP::VP6 and LT(R192G) on protection of BALB/c mouse against EDIM infection

Quantity of immunogen	#	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7	Shedding per mouse per day	% Reduction in shedding
0 $\mu$ g	1	0	11	862	1,190	208	79	0	385	
	2	0	363	3,491	592	508	62	0		
	3	0	220	2,491	846	184	116	4		
	4	0	85	943	490	184	221	24		
	5	0	62	740	584	110	146	0		
	6	0	41	2,480	1,248	652	151	0		
	7	0	6	242	663	118	108	0		
	8	0	20	217	532	137	117	4		
8.8 $\mu$ g	1	0	5	5	3	20	0	0	5	99
	2	0	0	8	0	0	0	0		
	3	0	29	0	0	0	0	0		
	4	0	16	5	0	0	0	0		
	5	0	13	8	0	0	0	0		
	6	0	68	26	3	0	0	0		
	7	0	14	10	0	0	0	0		
	8	0	7	0	0	23	0	0		
1.76 $\mu$ g	1	0	24	13	4	29	11	0	25	94
	2	6	32	80	0	0	3	0		
	3	0	40	74	3	15	5	0		
	4	0	143	18	0	116	0	0		
	5	0	47	113	6	48	6	0		
	6	0	72	9	19	92	17	0		
	7	0	21	25	0	22	0	0		
	8	6	76	61	0	134	0	0		

Although murine and human rotavirus VP6 proteins are highly homologous (see Example 30) it may be advantageous to have and use the human rotavirus VP6 protein for vaccination. Example 25 outlines the steps taken to clone the human rotavirus VP6.

#### Example 25

##### Expression of Human Rotavirus VP6

A VP6 protein from a human rotavirus strain is cloned and expressed as a fusion protein for development of a vaccine candidate to be tested in mice and humans. VP6 from human rotavirus strain CJN is cloned and its nucleotide sequence determined using standard techniques well known in the art. *See Current Protocols in Molecular Biology*, Eds. Ausubel, et al., John Wiley & Sons, Inc. The chimeric protein is tested in the mouse model to establish that a human VP6 protein from a group A virus can cross-protect against a heterologous group A (mouse EDIM) rotavirus. This human VP6 vaccine is tested in gnotobiotic pigs that have an immunological system similar to that of humans. This pig

model allows for the testing of whether mouse or human VP6 can protect against human rotaviruses. The protein is used in subsequent human trials.

Examples 26-29 provide outlines of vaccine trials in humans and various uses of the vaccine. Example 30 provides a way of testing which human haplotype(s) will be protective.

#### Example 26

##### Human Vaccine Trial

A statistically significant number of volunteers are enrolled in a study to test the safety and efficacy of the full-length, peptide, or chimeric rotavirus fusion protein vaccine compositions of the present invention. A geographical location or locations for the test is selected on the basis that the area is known to have been the site of past rotavirus outbreaks. The ratio of vaccine to placebo groups is randomized to result in a range from at least 1:1 to no more than 2:1 ratio within the group. This randomization is designed to provide appropriately large groups for statistical analysis of the efficacy of the vaccine.

The vaccine composition to be used in this study will be one containing the chimeric rotavirus fusion protein comprising VP6 and MBP, the full-length VP6 alone or in combination with a further rotavirus protein, or single or multiple peptide vaccines. The vaccine composition will consist of a sufficiently high concentration of rotavirus protein so as to be effective to induce a protective immune response when the composition is administered parenterally or mucosally. Parenteral administration will preferably be via intramuscular injection. In both cases any of the adjuvants which are disclosed in the specification can be used.

The chimeric fusion protein is prepared according to the Examples described above. The placebo will consist of an equal volume of buffered saline and is also to be given mucosally or parenterally. Vaccine and placebo are supplied as individual doses that are stored at -20°C and thawed immediately prior to use.

To determine the amount of vaccine necessary, different concentrations will be administered experimentally to a mouse. An effective concentration will be extrapolated and a comparable amount used in human subjects.

Blood samples are collected from all of the subjects for use in various laboratory assays. For example, enzyme immunoassay may be performed to evaluate the extent of the immune response elicited in each of the vaccinated individuals in response to the vaccine or placebo administered. Such techniques are well known in the art.

Individuals participating in this study are chosen who are healthy at the time of vaccination with either the test vaccine or the placebo. Test subjects are assigned to receive vaccine or placebo in a double-blind fashion using a block randomization scheme. An appropriate number of doses are administered over a given period of time, *e.g.*, two months, to elicit an immune response.

Study participants are monitored throughout the following year to determine the incidence of rotavirus infection and the subsequent development of disease conditions. Participating subjects are contacted on a periodic basis during this period to inquire about symptoms of rotaviral disease, both in the test subject and in the subject's community. Local epidemiological surveillance records may also be accessed.

The results of the above described study are assessed using standard statistical methods. The vaccine is well tolerated at the effective dose. The epidemic curves of outbreaks of rotavirus in the geographic areas tested will be assessed and the distribution of episodes of rotaviral disease will be established. The incidence of rotavirus caused disease in immunized individuals will be reduced to a statistically significant extent as compared to those individuals receiving the placebo.

#### Example 27

##### Maternal Immunization and Passive Immunity in Nursing Infants

In this example maternal immunization is used to induce a protective immune response in women of child bearing age which in turn passively protects nursing infants from rotavirus caused disease.

A woman of child bearing age is selected for immunization with the full-length, subunit, VP6 fragment, or VP6-fusion protein of the present invention. The woman is immunized parenterally.

Route of administration and the effective amount will be as discussed in Example 26.

#### Example 28

##### Infant and Maternal Immunization as a Method to Augment Infant Protection

Infants are immunized with a rotaviral protein of the present invention. The immunization is carried out by any route that results in the generation of a protective immune response directed against the rotaviral fusion protein immunogen, particularly intramuscularly. Route of administration and the effective amount will be as discussed in Example 26. Ideally, the resulting immune response will protect the infant from subsequent rotavirus challenge.

To augment the immunization of the infant, the mother or nursing female would also be immunized with the rotaviral protein vaccine of the present invention. Immunization is contemplated by any route that results in the generation of an immune response on the part of the immunized female, particularly intramuscularly. This immunization results in a protective immune response being generated on the part of the immunized mother. What is required is an immune response (soluble immunological factors) is induced against the rotaviral fusion protein. Those immunological factors should also be present in the milk generated by the immunized female during lactation. Accordingly, the nursing female may be immunized before nursing has begun.

The present invention contemplates a cooperative interaction between the immunized immune system of the infant and that of the immunized nursing female. In the interim between the time of infant immunization and the generation of an immune response, the passive transmission of immunity from the milk of the mother will protect the infant from rotavirus infection.

#### Example 29

##### Booster Immunization with a Subunit Vaccine to Augment Protection Obtained from Other Mono or Multivalent

##### Rotavirus Vaccines

In this Example an individual is first immunized with a traditional monovalent or multivalent rotavirus vaccine. Example of such vaccines may be found in U.S. Patent Nos. 4,927,628 and 5,626,851. Once an immune response has been mounted to this vaccine, rotaviral fusion proteins of the present invention may be used as a subsequent booster

to strengthen the immune response created by the first immunization. Route of administration and the effective amount will be as discussed in Example 26.

### Example 30

#### Human T cell proliferation studies

5 VP6 is extremely conserved among human rotavirus strains, and between human and animal rotaviruses (90-95% amino acid identity). In spite of the known HLA polymorphism, it is anticipated that human vaccines will generate protective immunity following i.n. vaccination with full-length VP6, possibly through responses to different CD4 epitopes which are conserved within multiple rotavirus strains and the degeneracy of T cell recognition of different HLA-peptide complexes. As already mentioned, BALB/c and Mt mice, which possess different H-2 molecules (H-2<sup>d</sup> and H-2<sup>b</sup>, respectively) on their T cells, are equally protected by MBP::VP6. To investigate whether humans with different haplotypes (HLA molecules) can be protected, one well-established approach to determine the presence of antigen-specific T cells is to perform proliferation assays. Such an assay may entail the following: blood is collected from a cross-section of subjects in a community and isolation of their peripheral blood mononuclear cells (PBMC) are isolated. PBMC are exposed *in vitro* to chimeric mouse or human MBP::VP6 or synthetic peptides. PMBC from individuals with the ability to generate VP6-specific T cell responses will proliferate. Proliferation of specific cells is detected by incorporation of <sup>3</sup>H-thymidine or using a non-radioactive proliferation assay (e.g. Molecular Probes, Eugene OR). Information from this experiment allows for further modification of the VP6 vaccine to ensure protection of the population targeted for vaccination.

SEQUENCE LISTING

	SEQ ID NO. 1	atg gat gtg ctg tac tct atc	Artificial Sequence
	SEQ ID NO. 2	tca cga gta gtc gaa tcc tgc aac	Artificial Sequence
5	SEQ ID NO. 3	atg gat gaa atg atg cga gag tca	Artificial Sequence
	SEQ ID NO. 4	tca gaa tgg cgg tct cat caa ttg	Artificial Sequence
	SEQ ID NO. 5	tgc gca att aat gct cca gct	Artificial Sequence
	SEQ ID NO. 6	tca ctt tac cag cat gct tct aat	Artificial Sequence
	SEQ ID NO. 7	atg gat gtg ctg tac tct atc	Artificial Sequence
10	SEQ ID NO. 8	tca gaa ctc aac ttc tac att att tgg	Artificial Sequence
	SEQ ID NO. 9	gca act aca tgg tac ttc aac cca	Artificial Sequence
	SEQ ID NO. 10	tca att tgg gaa aag tgc agt cac tgc	Artificial Sequence
	SEQ ID NO. 11	tca ttt caa ttg atg aga ccg cca	Artificial Sequence
	SEQ ID NO. 12	tca ttg tct gac tga cgt cac att ggc	Artificial Sequence
15	SEQ ID NO. 13	gaa tca gtt ctc gcg gat gca agt	Artificial Sequence
	SEQ ID NO. 14	tca ctt tac cag cat gct tct aat	Artificial Sequence
	SEQ ID NO. 15	CAINAPANIQQFEHIVQLRRVLTTA	Rotavirus VP6 fragment
	SEQ ID NO. 16	PDAERFSFPRVINSADGA	Rotavirus VP6 fragment
	SEQ ID NO. 17	FSFPRVINSADGATTWYFNPVILRPNNVEV	Rotavirus VP6 fragment
20	SEQ ID NO. 18	FNPVILRPNNVEVEFLNGQVINTYQARF	Rotavirus VP6 fragment
	SEQ ID NO. 19	NGQVINTYQARFGTIVARNFDTIRLSFQLM	Rotavirus VP6 fragment
	SEQ ID NO. 20	RNFDTIRLSFQLMRPPNMTPAVTAL	Rotavirus VP6 fragment
	SEQ ID NO. 21	MTPAVTALFPNAQPFHHATVGLTLRIDSA	Rotavirus VP6 fragment
	SEQ ID NO. 22	HATVLTLRIDSAICESVLADASETMLANV	Rotavirus VP6 fragment
25	SEQ ID NO. 23	VLADASETMLANVTSVRQEYAI	Rotavirus VP6 fragment
	SEQ ID NO. 24	QEYAI PVGPVFPFGMNWTDLITNYSPSRED	Rotavirus VP6 fragment
	SEQ ID NO. 25	TDLITNYSPSREDNLQRVFTVASIRSM LVK	Rotavirus VP6 fragment

WHAT IS CLAIMED IS:

1. A composition comprising a rotavirus VP6 protein or a fragment thereof, and an adjuvant in a pharmaceutical carrier, wherein said adjuvant is effective in combination with said VP6 protein or fragment thereof to generate a disease-reducing response to rotavirus infection in a mammal.
- 5 2. The composition of Claim 1, wherein said VP6 protein is derived from a human rotavirus strain.
3. The composition of Claim 1, further comprising an additional rotavirus protein selected from the group consisting of VP1, VP2, VP3, VP4, VP7, NSP1, NSP2, NSP3, NSP4, or NSP5.
4. The composition of Claim 1, wherein said adjuvant is selected from the group consisting of: cholera toxin (CT) and *E. coli* heat-labile toxin (LT).
- 10 5. The composition of Claim 1, wherein said adjuvant is selected from the group consisting of poly[di(carboxylatophenoxy)phosphazene] (PCPP), saponins Quil A, QS-7, and QS-21, the cholera toxin A1 subunit (CTA1) fused to a dimer of the Ig-binding D-region of *Staphylococcus aureus* protein A (DD) (CTA1-DD), and double-stranded RNA (dsRNA).
6. The composition of Claim 1, wherein said composition is adapted for administration by a route selected from the group consisting of intramuscular administration, intranasal administration, oral administration, transdermal administration, and transmucosal administration.
- 15 7. The composition of Claim 1, wherein said fragment is selected from the group consisting of; VP6<sub>AB</sub>, VP6<sub>BC</sub>, VP6<sub>CD</sub>, CD<sub>1</sub>, CD<sub>2</sub>, CD<sub>3</sub>, and CD<sub>4</sub>.
8. The composition of Claim 1, wherein said composition comprises a fragment of said VP6 protein, said fragment having a molecular weight less than about 24kD.
- 20 9. An isolated or purified peptide comprising a peptide with a sequence selected from the group of sequences consisting of SEQ ID NOs:15-24, and 25.
10. A recombinant rotavirus fusion protein composition, comprising:  
a rotavirus subunit fusion protein or fragment thereof, a fusion protein partner in genetic association with said recombinant rotavirus subunit protein, and an adjuvant in a pharmaceutical carrier, wherein said adjuvant is effective in stimulating a disease-reducing immunogenic response to said rotavirus fusion protein.
- 25 11. The composition of Claim 10, wherein said recombinant rotavirus fusion protein comprises a rotavirus subunit protein and a fusion partner protein in genetic association with said rotavirus subunit protein, wherein said fusion partner protein does not interfere with expression of said rotavirus subunit protein, said fusion partner protein prevents complex formation by said rotavirus subunit protein, and said fusion partner protein facilitates purification of said recombinant rotavirus fusion protein.
- 30 12. The composition of Claim 10, wherein said rotavirus subunit protein is selected from the group consisting of VP1, VP2, VP3, VP4, VP6, VP7, NSP1, NSP2, NSP3, NSP4, or NSP5.
13. The composition of Claim 12, wherein said rotavirus subunit protein is obtained from a human rotavirus strain.
- 35

14. The composition of Claim 10, wherein said fusion partner protein is selected from the group consisting of maltose binding protein, poly-histidine residues, S-Tag, glutathione-S-transferase, thioredoxin,  $\beta$ -galactosidase, nonapeptide epitope tag from influenza hemagglutinin, a 11-amino acid epitope tag from vesicular stomatitis virus, a 12-amino acid epitope from the heavy chain of human Protein C, green fluorescent protein, cholera holo toxin or its B subunit, labile holotoxin or its B subunit, streptavidin and dihydrofolate reductase.

15. The composition of Claim 10, wherein said fragment is selected from the group consisting of; VP6<sub>AB</sub>, VP6<sub>BC</sub>, VP6<sub>CD</sub>, CD<sub>1</sub>, CD<sub>2</sub>, CD<sub>3</sub>, and CD<sub>4</sub>.

16. The composition of Claim 10, wherein said adjuvant is selected from the group consisting of cholera toxin (CT), *E. coli* heat labile toxin (LT), poly[di(carboxylatophenoxy)phosphazene] (PCPP), saponins Quil A, QS-7, and QS-21, the cholera toxin A1 subunit (CTA1) fused to a dimer of the Ig-binding D-region of *Staphylococcus aureus* protein A (DD) (CTA1-DD), and double stranded RNA (dsRNA).

17. A full-length DNA copy of a gene encoding a recombinant rotavirus fusion protein, wherein said gene encoding said recombinant rotavirus protein comprising a rotavirus subunit protein or an immunogenic fragment thereof, and a fusion partner protein.

18. A host cell comprising the DNA clone of Claim 17.

19. A computer readable medium having recorded thereon the peptide sequence selected from the group of sequences consisting of SEQ ID NOs:15-24 and 25.

20. The computer readable medium of Claim 19, wherein said medium is selected from the group consisting of a floppy disk, a hard disk, random access memory (REM), read only memory (ROM), and CD-ROM.

21. Use of an immunogenic composition according to any one of the forgoing claims for the manufacture of a medicament for generating a rotaviral disease-reducing immunogenic response.

22. The use of Claim 21, wherein the rotavirus protein is VP6.

23. The use of Claim 22, wherein the rotavirus VP6 protein is obtained from a human rotavirus strain.


24. The use of Claim 21 wherein said rotavirus protein is a fusion protein.

25. The use of Claim 21, wherein the composition is adapted for administration by a route selected from the group consisting of intramuscular administration, intranasal administration, oral administration, transdermal administration, and transmucosal administration.

26. The use of any one of claims 1-16 in the preparation of a medicament for boosting an immune response to a previously administered rotavirus vaccine.



Figure 1. Plasmid map of pMAL-c2 used for expression of chimeric MBP proteins genetically fused to rotavirus proteins



**Figure 2. Western blot analyses of chimeric MBP proteins genetically fused with rotavirus VP4, VP6 or a truncated VP7 protein**




Figure 3. Western blot analyses of MBP::VP6 and silver stained gel of rotavirus particles in fractions collected after ultracentrifugation through a sucrose gradient.




FIGURE 4

Intranasal- VP6 + LT




FIGURE 5

Regions of VP6 cloned into pMAL-c2. VP6 is delineated into regions A, B, C and D. pMAL-c2/EDIM6<sub>AB</sub> encodes regions A and B, pMAL-c2/EDIM6<sub>BC</sub> encodes regions B and C, and pMAL-c2/EDIM6<sub>CD</sub> encodes regions C and D.

Plasmid	Amino acids encoded	Regions of VP6			
		A	B	C	D
PMAL-c2/EDIM	1-397	— — — —			
PMAL-c2/EDIM6 <sub>AB</sub>	1-196	— —			
PMAL-c2/EDIM6 <sub>BC</sub>	97-299		— —		
PMAL-c2/EDIM6 <sub>CD</sub>	197-397			— —	

6/7

Figure 6. Western blot analyses of antisera collected from mice immunized with plasmids expressing chimeric MBP genetically fused with VP<sub>6AB</sub>, VP<sub>6BC</sub>, or VP<sub>6CD</sub>.









FIGURE 7

Regions of VP6<sub>CD</sub> cloned into pMAL-c2. The CD region of VP6 is delineated into subregions 1, 2, 3 and 4. These subregions were cloned into pMAL-c2 to create pMAL-c2/EDIM6<sub>CD1</sub>, pMAL-c2/EDIM6<sub>CD2</sub>, pMAL-c2/EDIM6<sub>CD3</sub> and pMAL-c2/EDIM6<sub>CD4</sub>.

Plasmids	Amino acids encoded	Regions of VP6
pMAL-c2/EDIM6 <sub>CD</sub>	197-397	
pMAL-c2/EDIM6 <sub>CD1</sub>	197-263	
pMAL-c2/EDIM6 <sub>CD2</sub>	244-310	
pMAL-c2/EDIM6 <sub>CD3</sub>	291-351	
pMAL-c2/EDIM6 <sub>CD4</sub>	332-397	

## INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 99/25799

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/46 C12N15/62 C07K14/14 C07K19/00 A61K39/15

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	O'NEAL C. M. ET AL.: "ROTAVIRUS 2/6 VIRUSLIKE PARTICLES ADMINISTERED INTRANASALLY WITH CHOLERA TOXIN ESCHERICHIA COLI HEAT-LABILE TOXIN (LT) AND LT-R192G INDUCE PROTECTION FROM ROTAVIRUS CHALLENGE" JOURNAL OF VIROLOGY, vol. 72, no. 4, April 1998 (1998-04), pages 3390-3393, XP002133270	1-4, 6, 21-23, 25, 26
Y	the whole document	5, 7-16, 24
	— — — — — — / —	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance  
 "E" earlier document but published on or after the international filing date  
 "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)  
 "O" document referring to an oral disclosure, use, exhibition or other means  
 "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention  
 "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone  
 "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art  
 "&" document member of the same patent family

Date of the actual completion of the international search

16 March 2000

Date of mailing of the international search report

04/04/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax: (+31-70) 340-3016

Authorized officer

Mandl, B



## INTERNATIONAL SEARCH REPORT

 Internat Application No  
 PCT/US 99/25799

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	MCNEAL M. M. ET AL.: "STIMULATION OF LOCAL IMMUNITY AND PROTECTION IN MICE BY INTRAMUSCULAR IMMUNIZATION WITH TRIPLE- OR DOUBLE-LAYERED ROTAVIRUS PARTICLES AND QS-21" VIROLOGY, vol. 243, no. 1, 30 March 1998 (1998-03-30), pages 158-166, XP002133271 the whole document	5
X	BAÑOS D. M. ET AL.: "Identification of a T-helper cell epitope on the rotavirus VP6 protein." JOURNAL OF VIROLOGY, vol. 71, no. 1, January 1997 (1997-01), pages 419-426, XP002133272 the whole document	9,17-20
Y		7-16,24
X	EP 0 235 754 A (ABBOTT LAB) 9 September 1987 (1987-09-09)  the whole document	1,2,6,8, 10-14, 17,18, 21-26
X	WO 92 07941 A (UNIV SASKATCHEWAN) 14 May 1992 (1992-05-14) the whole document	1-3,6, 20-26
X	EP 0 235 391 A (UNIV SASKATCHEWAN) 9 September 1987 (1987-09-09)  page 10, line 35 -page 12, line 21 page 29, line 34 -page 30, line 27 table 7	1-3,6,8, 10-13, 21-26
A	KOHLI E. ET AL.: "Epitope mapping of the major inner capsid protein of group A rotavirus using peptide synthesis." VIROLOGY, vol. 194, 1993, pages 110-116, XP002133273 the whole document	1-26
P,X	CHOI A. H. ET AL.: "ANTIBODY-INDEPENDENT PROTECTION AGAINST ROTAVIRUS INFECTION OF MICE STIMULATED BY INTRANASAL IMMUNIZATION WITH CHIMERIC VP4 OR VP6 PROTEIN" JOURNAL OF VIROLOGY, vol. 73, no. 9, September 1999 (1999-09), pages 7574-7581, XP002133274 the whole document	1,4-6, 10-12, 14, 16-18, 21,22, 24-26

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 99/ 25799

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

See additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☒ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest.

☐ No protest accompanied the payment of additional search fees.

**FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210**

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-8 (completely), 21-23,25,26 (partially)

A composition comprising a rotavirus VP6 protein or a fragment thereof and an adjuvant; and the use of said composition for the manufacture of a medicament.

2. Claims: 9,19,20 (completely), 21-23,25,26 (partially)

An isolated or purified peptide selected from the group of peptides represented by SEQ.ID.s 15-25; a computer readable medium having recorded thereon said peptide sequences; and the use of said peptide for the manufacture of a medicament.

3. Claims: 10-18,24 (completely), 21-23,25,26 (partially)

A recombinant rotavirus fusion protein composition comprising a rotavirus subunit fusion protein and an adjuvant; a DNA molecule encoding said fusion protein; and the use of said composition for the manufacture of a medicament.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US 99/25799

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0235754 A	09-09-1987	JP 62234098 A	14-10-1987
W0 9207941 A	14-05-1992	US 5298244 A	29-03-1994
		AT 152768 T	15-05-1997
		AU 659416 B	18-05-1995
		AU 8718791 A	26-05-1992
		CA 2094916 A	26-04-1992
		DE 69126043 D	12-06-1997
		DE 69126043 T	18-09-1997
		DK 652957 T	27-10-1997
		EP 0652957 A	17-05-1995
		ES 2103828 T	01-10-1997
		GR 3024173 T	31-10-1997
		JP 6504190 T	19-05-1994
EP 0235391 A	09-09-1987	AU 600460 B	16-08-1990
		AU 6698786 A	02-07-1987
		CA 1327096 A	15-02-1994
		CN 1020857 B	26-05-1993
		DE 3685651 A	16-07-1992
		JP 2616915 B	04-06-1997
		JP 63115825 A	20-05-1988